



Review Article

Anestrus in Cattle and Buffalo: Indian Perspective

Priya Ranjan Kumar^{1*}, Sanjay Kumar Singh¹, Suresh Dinkar Kharche², Chethan Sharma Govindaraju¹, Bijay Kumar Behera¹, Satya Nidhi Shukla³, Harendra Kumar¹, Sudhir Kumar Agarwal²

¹Division of Animal Reproduction, Indian Veterinary Research Institute, Izatnagar, Uttar Pradesh- 243122 (India); ²Central Institute for Research on Goats, Makhdoom, Mathura-281122 (India), ³Department of Veterinary Gynaecology and Obstetrics, College of Veterinary Science and Animal Husbandry, Nanaji Deshmukh Veterinary Science University, Jabalpur Madhya Pradesh-482001 (India)

*Corresponding author: dr.pranjan007@gmail.com

ARTICLE HISTORY

Received: 2014-01-18
Revised: 2014-01-21
Accepted: 2014-01-22

Key Words: Anestrus, Cattle, Buffalo, Causes, Diagnosis, Treatment

ABSTRACT

Anestrus is a functional disorder of reproductive cycle in cattle and buffalo which is characterized by absence of overt signs of estrus and also affecting the livestock enterprise to a great extent. Incidence of anestrus is more in buffalo than the cattle, and problem is severe during summer. Anestrus is a multifactorial problem but its occurrence signals the inadequate nutrition, environmental stress, uterine pathology and improper managemental practices. It can be classified based on physiological and pathological conditions of the animals. Diagnosis is based on the exploration of the different causative factors involved in. Many therapeutic agents (hormonal and non hormonal) have been used, however, as such there is no single panacea to correct it. Though, recruitment and emergence of follicular wave is usually not affected, the derangement in subsequent growth and development and ovulation of dominant follicle leads to anovulatory anestrus. In order to achieve efficient management strategies, further research is needed for better understanding of its etiology, diagnosis and therapeutics.

All copyrights reserved to Nexus® academic publishers

ARTICLE CITATION: Kumar PR, Singh SK, Kharche SD, Chethan Sharma G, Behera BK, Shukla SN, Kumar H, Agarwal SK (2014). Anestrus in cattle and buffalo: Indian perspective. *Adv. Anim. Vet. Sci.* 2 (3): 124 – 138.

INTRODUCTION

Anestrus is one of the most commonly occurring reproductive problems in cattle and buffalo in India, affecting livestock productivity and economics to a great extent. The problem is more severe in sub urban and rural areas of the country. It is a functional disorder of the reproductive cycle which is characterized by absence of overt signs of estrus manifested either due to lack of expression of estrus or failure of its detection. Anestrus is observed in post pubertal heifers, during pregnancy, lactation and in early postpartum period in adult animals. The condition may be associated with uterine pathology such as pyometra, fetal resorption, maceration and mummification. Expression of estrus is also influenced by seasonal changes, stress and aging. In heifers, it poses a herd problem possibly due to low plane of nutrition, stress of seasonal transition or extremes of climatic conditions. Expression of overt signs of estrus is greatly affected by heat stress in buffaloes. Modern feeding and managemental practices also accentuate the problem in commercial dairy farms. Incidence of anestrus though varies in the different managemental system but it is more in buffalo than the cattle, and especially during summer. Anestrus is a multi-causative factors associated problem but its occurrence signals the inadequate nutrition, environmental stress, uterine pathology and improper managemental practices. Diagnosis of the condition is based on the exploration of the different causative factors responsible for it. Though many

therapeutic agents (hormonal and non-hormonal) have been used but as such there is no single panacea to correct it. The present review on “Anestrus in cattle and buffalo” has been described in details keeping in mind the Indian perspective.

INCIDENCE

A large variation on incidence of anestrus has been reported in literatures depending upon species, breed, parity, season, level of nutrition, managemental conditions, geographic environment (Table 1 and 2). In general, incidence in India has been reported between 2.13–67.11 and 9.09–82.50 per cent in indigenous cattle (Dessouky and Juma, 1973; Kodagali, 1974; Luktuke and Sharma, 1978; Pargaonkar, 1978; Singh et al., 1981; Rehman et al., 1990; Iyer et al., 1992; Narladkar et al., 1994; Singh, 2003; Yadav et al., 2004; Selvaraju et al., 2005; Thakor and Patel, 2013) and buffaloes (Kodagali, 1968; Luktuke et al., 1973; Singh and Verma, 1994; Nanda et al., 2003; Prajapati et al., 2005; Singh et al. 2006; Modi et al., 2011; Kumar et al., 2013; Thakor and Patel, 2013), respectively. The incidence of anestrus among crossbred cattle has been reported between 2.55–40.4 per cent (Dessouky and Juma, 1973; Narladkar et al., 1994) from different parts of the country. Its incidence in heifers has been reported between 12.37 to 64.66 per cent (Luktuke and Sharma, 1978; Naidu and Rao, 1981; Nayak and Mohanty, 1986; Sinha et al., 1987). Incidence of anestrus is higher in adult cattle and buffalo than the heifers (Bharkad and Markandeya, 2003).

Table 1: Incidence of Anestrus in cattle and buffalo in India state-wise pattern

States	Cattle Incidence (%)	References	Buffalo Incidence (%)	References
Andhra Pradesh	49.70	Rao, 1993	30.76–50	Rao and Sreemannarayanan, 1982; Naidu, 2004
Gujarat	24.73	Patel et al., 2007	20.84 – 45.97	Kulkarni, 2002; Modi et al., 2011
Madhya Pradesh	53.15	Pandit, 2004	29.12 – 60. 83	Pandit, 2004; Kumar et al., 2013
Maharashtra	2.13–45.97	Narladkar et al., 1994; Kulkarni et al., 2002	29.5–41.40	Bharkad and Markandeya, 2003;
Tamilnadu	16.6	Selvaraju et al., 2005	9.09	Selvaraju et al., 2005
Punjab	43–67.11	Singh et al., 2003	38.98–55.5	Singh et al., 2006
Bihar	39.01	Singh, 1981	–	–
Kerala	65.0	Kutty and Ramachandran 2003	–	–
Kashmir	27.52 & 31.0	Rehman et al., 1990; Bhattacharyya and Buchoo, 2008	–	–
Utter Pradesh	–	–	14.69–45.20	Luktuke et al., 1973
Karnataka	–	–	56.0	Hussain, 1984

Table 2: Incidence of anestrus in India- breeds wise

Breed	Incidence (%)	References
Hariyana cows	0.6–12.13	Khan and Luktuke, 1967
Kankrej cows	11.8–23.8	Agarwal and Buck, 1968
Gir cows	47.89	Kodagali, 1974
Red Kandhari cows	3.0	Pargaonkar and Bakshi, 1987
Deoni cows	2.13	Narladkar et al., 1994
Friesian crossbred	35.96	Sinha et al., 1987
Jersey crossbred	35.5	Sinha et al., 1987
Murrah and upgraded Murrah buffaloes	29.12	Kumar et al., 2013

The period of postpartum anoestrus is usually longer in buffalo than the cattle under similar management conditions (Dobson and Kamonpatana, 1986; Jainudeen and Hafez, 1993). In comparison to cows, buffaloes have lesser number of preantral and antral follicles, smaller sized pre-ovulatory follicle and greater tendency of follicular atresia (Samad and Nasser, 1979; Danell, 1987; Baruselli et al., 1997) which might be responsible for high incidence of anestrus in buffaloes.

ECONOMIC IMPACT

High milk production and excellent fertility are desirable traits for a profitable dairy enterprise. Infertility due to cyclicity failure or anestrus has great economic impact. Anestrus, leads to economic losses through increased intercalving interval, poor net calf crops, production loss, treatment expenses and cost of replacing mature animal with first calving heifer. There are only few reports from India pertaining to economic impact analysis due to anestrus. Pawshe et al. (2011) reported an estimated loss from anestrus around Rs.193.00 per day in cow and Kumar et al. (2013) reported Rs. 372.90 per day in buffalo. As the incidence of anestrus in India has been reported high, the above figures show great economic impact at country level.

CLASSIFICATION OF ANESTRUS

Based on ovarian activity, anestrus cow has broadly been classified into ovulatory, anovulatory and inactive. Robert (1971) divided the anestrus cow or heifer into two class viz., Class I-cow with a normal functional corpus luteum and Class II-cow with no functional corpus luteum. Wiltbank

et al. (2002) classified the anovulatory anestrus into three classes i.e. anovulation with follicular growth up to emergence; anovulation with follicular growth up to deviation but not ovulatory one; anovulation with follicular growth up to ovulatory size. Further, Peter et al. (2009) divided anestrus into four types viz., Type-I, characterized by emergence of follicular wave and growth of follicle up to pre deviation stage; Type-II, characterized by deviation of follicle and follicle grow up to dominance stage followed by atresia; Type-III, characterized by persistent dominant follicle (persistent of large follicle) i.e. follicle grow up to preovulatory or ovulatory size or more but fails to ovulate or regress (follicular and luteal cyst) and Type-IV, characterized by normal ovulation and formation of corpus luteum but corpus luteum persist beyond the expected time of regression (persistent corpus luteum, PCL) resulting into anestrus.

Based on the above informations and for the ease, anestrus has been classified accordingly for better understanding (Figure 1).

PHYSIOLOGICAL ANESTRUS

Animals remain anestrus during certain physiological stages which does not related to infertility viz., before puberty, during pregnancy, lactation and early postpartum period. Accordingly, physiological anestrus has been classified into pre-pubertal, gestational, lactational and post-partum anestrus.

PREPUBERTAL ANESTRUS

The follicular waves in pre-pubertal animals are similar to that of adult but follicles grow in response to FSH secretion only up to the stage where they have a theca interna and then regress. Such heifers remain in anestrus before the onset of puberty. The reasons of pre-pubertal anestrus includes low LH pulse frequency that results in insufficient growth of follicles; inhibitory effect of opioids on LH secretion and high threshold for positive feedback effect of estradiol on LH surge (Noakes et al., 2009).

GESTATIONAL ANESTRUS

The elevated level of progesterone during pregnancy exerts negative feedback effect on GnRH secretion from hypothalamus and reduces LH pulse frequency resulting into anestrus. However, some cattle and buffaloes exhibit estrus during early pregnancy (known as gestational estrus) which is seen most often during first four months of pregnancy. The incidence of gestational estrus has been recorded as 3.33 to 20.3% and 6.05 to 14.40% in Indian cattle (Luktuke et al., 1964; Chauhan et al., 1976; Kaikini and Fasihuddin, 1984) and buffaloes (Luktuke et al., 1964),

respectively. Usually cow or buffalo exhibits gestational heat only once during pregnancy, however, few animals show twice or thrice in same gestation.

POSTPARTUM ANESTRUS

Following parturition, all the females undergo through anestrus for a variable but short period of time, known as postpartum anestrus. The period of postpartum anoestrus is usually longer in buffalo than the cattle under similar management conditions (Dobson and Kamonpatana, 1986; Jainudeen and Hafez, 1993), probably due to low LH secretion during early postpartum period (Perera, 2011). Under normal conditions, buffaloes resume cyclicity by 30–90 days (Perera, 2011), however; only about 45% of Indian buffaloes resume cyclicity within 90 days postpartum and rest 55% remain in anestrus for about 150 days (El-Wishy, 2007). Most of the dairy cows resume ovulatory estrus cycle within 15–45 days postpartum (Butler and Smith, 1989; Forde et al., 2011). The physiological postpartum anestrus cannot be avoided and is useful to allow uterine involution prior to first postpartum anestrus.




Figure 1: Classification of anestrus in cattle and buffalo

LACTATIONAL ANESTRUS

High lactation suppresses the fertility in almost all the mammals (Warnick et al., 1950; Baker et al., 1953). Higher level of prolactin in high yielding animals suppresses GnRH secretion and ultimately reduces production of gonadotrophins from pituitary, resulting into anestrus.

PATHOLOGICAL CAUSES OF ANESTRUS

Certain pathological conditions i.e. ovarian agenesis, dysgenesis or derangement of follicular-luteal dynamics leads to anestrus causing infertility and pose a herd problem. Such conditions may be congenital or acquired.

CONGENITAL AND HEREDITARY CAUSES OF ANESTRUS

Congenital and hereditary form of anestrus is upshot of ovarian agenesis or dysgenesis. Ovarian agenesis or aplasia (absence of ovary) is extremely rare condition and probably crop up due to inherited autosomal dominant gene. Bilateral aplastic or gonadless heifers appear normal until breeding age but fail to show estrus and normal development of udder at puberty and are sterile. Such reports from India are meager. Ovarian dysgenesis has been identified as ovarian hypoplasia and freemartin. Ovarian hypoplasia (incomplete development of ovary) is caused by single autosomal recessive gene with incomplete penetration. It may be unilateral or bilateral. The affected ovary is characterized by lack of primordial follicles reserve either partial (partial hypoplasia) or complete (complete hypoplasia) (Settergren, 1964, 1997). Bilateral complete hypoplastic females remain in anestrus whereas partial hypoplastic animals exhibit estrus, conceive and produce viable calves but transmit this undesirable character to the next generation therefore must be avoided. The small size of normal ovary in buffalo heifers often mistaken as hypoplastic ovaries. The incidence of ovarian hypoplasia in Indian cattle has been reported between 0.08–4.3% (Kodagali, 1969; Narasimha Rao and Murthy, 1972; Nair and Raja, 1974; Bonia, 1981; Kumar and Agarwal, 1986) whereas it is less than 1 per cent in Indian buffaloes (Damodaran, 1956; Malik et al., 1960; Narasimha Rao and Sreemannarayana, 1982; Mittal et al., 2010) with a slightly high incidence (1.46%) in Jaffarabadi buffalo (Kodagali and Kerur, 1968). Higher incidence of ovarian hypoplasia (10–23%) has been reported in exotic cattle (Lagerlof and Boyd, 1953). In Free martin (sterile heifer born co-twin with bull calf) the ovaries usually fails to develop and remain hypoplastic resulting into anestrus. The incidence of freemartinism in Indian cattle and buffaloes has been reported low i.e. between 0.10 to 0.20% (Bhagat, 1966; Narasimha Rao and Suryanarayana Murthy 1980, Sharma et al., 2004).

ANOVULATORY ANESTRUS-I

This condition is characterized by growth of follicles only to a stage of follicular wave emergence i.e. up to about 4 mm diameter and then regress (Wiltbank et al., 2002). This type of anestrus is very rare and only 0.003% cases has been reported by Wiltbank et al. (2002), however, expected to occur in about 10% of postpartum dairy cows (Zemjanis et al., 1969; Markusfeld, 1987). Ovaries of such animals are small and smooth with absence of corpus luteum and classified as “inactive ovaries” in earlier literatures. The

factors responsible for growth of follicles up to this stage are poorly known, however, to be sure that these events are gonadotrophin independent. On the basis of *in-vitro* studies, various peptide growth factors and cytokines have been implicated as positive (e.g. Kit Ligand, insulin, LIF, FGF2, BMP4, BMP7 and GDF9) and negative (e.g. AMH and CXCL12) regulators of follicular growth at this stage (Kezele et al., 2002; Fortune, 2003; McLaughlin and McIver, 2009). The occurrence of anovulatory anestrus-I is possibly due to deficiency of one or more of these growth regulators presumed to occurs under extreme undernutrition or severe energy deficit.

ANOVULATORY ANESTRUS-II

This type of anestrus animals exhibit normal follicular wave pattern and follicular growth proceeds through emergence but develop only up to deviation or preovulatory stage (not ovulatory size) followed by regression and emergence of new follicular wave 2–3 days later (Roche et al., 1998; Wiltbank et al., 2002; Peter et al., 2009; Ghuman et al., 2010). This process of follicular growth and regression repeated over and again in anestrus animals. It is most common form of anestrus. The ovaries of such animals are small with the absence of a corpus luteum or ovulatory size follicles. Adequate LH pulse frequency is required for growth and development of follicle after emergence of follicular wave. The occurrence of this condition is due to low LH pulse frequency (1/3–4hrs).

ANOVULATORY ANESTRUS-III

In this type of anestrus, follicular growth proceeds through emergence and deviation stage and attain dominance but fails to ovulate and become persistent. Persistent follicle may develop either into follicular cyst or luteal cyst. Follicular cysts are thin walled structure with no evidence of luteinization of granulosa cells (progesterone level <1ng/ml) and characterized by either anestrus or nymphomania (persistent estrus), whereas luteal cysts are thick walled structures with partial luteinization of granulosa cells and characterized by anestrus. This type of anestrus is up shot of absence or attenuation of LH surge (Cook et al., 1990; Hamilton et al., 1995, Peter et al., 2004) due to partial or complete failure of estradiol to bring out normal LH surge (Nanda et al., 1991). This appears to be due to insensitivity of hypothalamic surge centre to the positive feedback of estradiol (Garverick, 2007) and failure of GnRH release (Vanholder et al., 2005). Another probable causes for the development of cysts are occurrence of LH surge at an inappropriate time i.e. when follicle is not capable to ovulate (Vanholder et al., 2005) or reduced population of LH receptors on follicle (Kawate, 2004). In India, the incidence of follicular cysts has been reported 0.5–2.7% in cattle (Kulkarni et al., 2002) and 1.26–3.4% in buffaloes (Bhattacharya et al., 1954; Malik et al., 1960) and 8.0–10.2% in crossbred cattle, whereas incidence of luteal cysts are 1.45–3.0% in cattle (Luktuke and Arora, 1967; Singh, 1981; Kulkarni et al., 2002), 0.66–12.5% in buffaloes (Damodaran, 1956; Rao 1965; Kaikini, 1974; Kuty and Ramachandran, 2003) and 9.0–13.3% in crossbred cattle (Kulkarni et al., 2002). The incidence of follicular and luteal cysts abroad in dairy cattle have been reported 6–19% (Garverick, 1997) and 10–13% (Peter, 2004), respectively, however, these are less frequent in dairy heifers and beef cattle.

ANESTRUS DUE TO PERSISTENT CORPUS LUTEUM (PCL)

In this type of anestrus, the follicular growth proceeds through all the developmental stages and undergo ovulation and CL formation which subsequently turn into anestrus due to failure of luteal regression. This is probably due to absence of estrogenic dominant follicle at the time of luteal regression (Wiltbank et al., 2002) secreting adequate estradiol to induce the formation of uterine oxytocin receptors and consequently resulting in to pulsatile release of PGF₂ α for luteolysis (Knickerbocker et al., 1986; Thatcher et al., 1989; McCracken et al., 1999). Persistent corpus luteum (PCL) is mostly associated with uterine pathology such as endometritis, pyometra, fetal resorption, maceration, mummification and uterine unicornis (Lynn et al., 1966; Gilbert et al., 1990; Noakes et al., 1990). Retained corpus luteum may also be associated with embryonic death when death of embryo occurs after maternal recognition of pregnancy where corpus luteum persists until resorption of embryo. In uterine unicornis, CL formed on the ovary ipsilateral to the missing uterine horn persists due to a lack of a luteolytic signal from the missing uterine horn. PCL results in anestrus due to inhibitory effect of progesterone secreted by the corpus luteum on hypothalamo-hypophyseal axis for the secretion of gonadotrophins.

SUB-ESTRUS/SILENT ESTRUS/QUITE OVULATION

Sub estrus or silent estrus or quite ovulation is clinically characterized by failure of overt symptoms of estrus, though the animal is surprisingly normal. Under these conditions, follicular development and ovulation occurs normally in animals without the manifestation of overt signs of estrus. Sub estrus is common during the post pubertal period in heifers and early post-partum (30 to 120 days) in high yielding dairy cows. Progesterone secreted from regressing CL of previous cycle potentiates the action of estrogen and seems to favours the manifestation of estrus in next cycle (King et al., 1976; Allrich, 1994). Thus, lack of progesterone priming results in sub-estrus. Such conditions have been frequently reported in dairy buffaloes especially in summer months (Sane et al., 1967; Shah, 1990; Badr, 1993; Singh et al., 2013) and may be the one of the reasons of prolonged calving interval in buffaloes (Singhal et al., 1984; Barkawi et al., 1986). The concentration of estrogen determines intensity of behavioral signs of estrus which is low in high yielding dairy cow (Lopez et al., 2004). Lower concentration of estrogen may be either due to higher metabolism and clearance with a high metabolic load (Sangsritavong et al., 2002) or sub-optimal follicular growth (Awasthi et al., 2007). The probable cause of silent estrus is sub-optimal secretion of estradiol by mature follicles or higher threshold of estrogen in central nervous system to display the symptoms of estrus in that particular individual animal. Silent estrus also appears to be hereditary in predisposition in certain breeds (Lagerlof, 1951). Other causes of sub estrus are heat stress, nutritional deficiencies, overweight, foot lesions, aging and ergotism (fescue toxicity) but most common cause considered for sub estrus is the failure of estrus detection.

ANESTRUS DUE TO FAILURE TO OBSERVE ESTRUS/UNOBSERVED ESTRUS

Estrus detection is critical aspect of dairy herd management where artificial insemination is being practiced. The length

of estrous cycle and estrus period varies among breeds and within a breed (Rottensten and Touchberry, 1957; Roxström, et al., 2001). It also varies with season, nutrition, lameness, presence of bull, housing, herd size and production status (King et al., 1976; Purbey and Sane, 1978; Zakari et al., 1981; Lucy, 2003; Lopez et al., 2004; Diskin, 2008; Walker et al., 2008). Earlier, it was reported that intensity and duration of standing estrus is shorter in *Bos indicus* cattle as compared to *Bos taurus* cattle (Anderson, 1936; De Alba et al., 1961; Plasse et al., 1970; Rhodes and Randel, 1978), probably due to small follicular diameter (Lyimo et al., 2000; Bo et al., 2003). However, recent studies indicate that there is no difference in intensity and duration of estrus between *Bos taurus* and *Bos indicus* cows (Bastos et al., 2010). In high yielding cows, many times the estrus cycles become irregular (Bartha, 1971) in terms of its intensity and duration of standing estrus (Lopez et al., 2004), resulting in low estrus detection rates. The condition may be due to low estrogen concentration (Lyimo et al., 2000; Lopez et al., 2004), insulin and IGF-I mediated deficiency of follicular growth (Butler and Smith, 1989; Lucy, 2001) or increase metabolism and clearance of estrogen with high metabolic load (Sangsritavong et al., 2002). The short period of estrus often fail to notice by the farmers.

PATHOGENESIS

The concentration of gonadotrophins are almost negligible in late gestation and for a short duration following parturition due to strong inhibition of hypothalamic-pituitary axis through negative feedback effect of high progesterone secreted by corpus luteum and placenta and estrogen from placenta during last trimester of pregnancy. The concentration of FSH rises within 3–5 (ranges 2 to 7) days after parturition (Ginther et al., 1996; Crowe et al., 1998) whereas restoration of LH pulsatility and LH surge mechanism takes rather longer period than the FSH (14–28 days). Emergence of follicular wave occurs within 1 or 2 days of significant rise in FSH concentration (Beam and Butler, 1997; Crowe et al., 1998), however, none of the growing follicles become mature enough to ovulate due to very low LH pulse frequency (Canfield and Butler, 1990; Montiel and Ahuja, 2005; Roche, 2006). Moreover, growth and maturation of follicle also depends upon bioavailability of insulin, insulin like growth factor (IGF)-I and their binding proteins (Butler and Smith, 1989; Lucy, 2001; Fortune et al., 2004). Both insulin and IGF-I are potent stimulators of steroidogenesis and granulosa and theca cells proliferation as well as oocytes growth and maturation (Webb et al., 1999; Lucy et al., 1999; Silva et al., 2009). Hence, inadequate LH pulse frequency, low concentration of insulin and IGF-I impede the follicular growth and reduces the chance of ovulation. These conditions appear to occur in a state of under nutrition/malnutrition (Roche, 2006; Ramoun et al., 2012) and negative energy balance (Richards et al., 1989; Beever et al., 2001). In a state of negative energy balance, the circulating concentration of non-esterified fatty acid (NEFA) due to mobilization of body reserves (Richards et al., 1989; Canfield and Butler, 1990; Grimard et al., 1995) and endogenous opioids increases which in turn decreases the pulsatile secretion of LH. In addition to these, prolactin, oxytocin, cortisol, corticotrophin releasing hormone secreted in response to

lactation, suckling and stress also reduces the LH pulse frequency and subsequently impair the follicular growth. Ultimately, follicles become atretic and then regress. The process of follicular growth and regression occurs over and again till the above state of affairs persists resulting into anovulatory anestrus.

FACTORS AFFECTING ANESTRUS

Nutrition

Nutritional status of animals affects the follicular growth, maturation and ovulation (Diskin et al., 2003). Under nutrition is the one of the most prevalent cause of anestrus in heifers. Extended postpartum period of anestrus (>150 days) are usually observed in cattle of tropical area under free range rearing system, probably due to shortage of feed and good quality fodder. Reduced feed intake during late gestation or/and early postpartum period or negative energy balance (NEB) due to very high metabolic load following parturition especially in high yielders delays postpartum restoration of LH pulsatility, resulting into prolonged postpartum anestrus (Wiltbank et al., 1962; Rutter and Randel, 1984; Easdon et al., 1985; Nolan et al., 1988; Connor et al., 1990; Hegazi et al., 1994). Under high metabolic load, nutrients are utilized for production rather than reproduction (Ferguson, 2001). In addition to NEB, the deficiency of minerals like calcium (Ca), phosphorus (P), copper (Cu), zinc (Zn) and manganese (Mn) are also associated with anestrus (Hidiroglou, 1979; Campbell et al., 1999). It is well established that minerals play an intermediate role in the action of hormones and enzymes at cellular level and its deficiency ultimately affect the reproductive performance of females (Bearden et al., 2004).

Occurrence of anestrus is not only due to underfeeding or malnutrition but also occurs owing to high feed intake which promotes high metabolism and clearance of ovarian steroids (estrogen and progesterone) from body by enhancing the hepatic perfusion (Sangsrivong et al., 2002), especially in high yielders. Low estrogen in high yielding animals result in sub estrus as intensity of estrus behaviors is directly related to its concentration (Lymio et al., 2000; Lopez et al., 2004). Excessive clearance of ovarian steroids also leads to anovulation (Walsh et al., 2007) and delayed luteal regression (Opsomer et al., 2000; Petersson et al., 2006).

Body Condition Score (BCS)

Body condition score is the measures of nutritional status of animals and is an important factor influencing the reproductive performance (Baruselli et al., 2001). Extremes of BCS (very low and very high) at pre-calving, calving and early postpartum period delay onset of cyclicity (Butler and Smith, 1989; Markusfeld et al., 1997; Pryce et al., 2000). However, BCS at calving is a better indicator of resumption of postpartum cyclicity than prepartum BCS (Whitman, 1975; Lalman et al., 1997). For optimum reproductive performance, BCS of 3.5 (on five point scale) is required at calving (Bhalaru et al., 1987; Ribeiro et al., 1997). Restricted feed intake during late gestation and early postpartum period result in low BCS, consequently, leads to prolonged postpartum anestrus (Dziuk and Bellows, 1983; Robinson, 1990).

Environmental Stress

Environmental stress (extreme cold and heat) affect the development of follicles and manifestation of estrus both in

cattle and buffaloes. Although buffaloes are well adopted in hot and humid climate, however, ovarian activity is greatly reduced by heat stress and manifested in the form of anestrus (Singh et al., 2000). Heat stress affects folliculogenesis, follicular fluid micro environment and oocyte quality. In buffaloes, decline in feed intake during summer results in reduced secretion of gonadotrophins (El-Sawaf et al., 1979). Besides this, high environmental temperature causes hyper-prolactinaemia and suppressing the secretion of gonadotrophin which leads to alteration in ovarian folliculogenesis and steroidogenesis. Among cattle, reproductive functions are suppressed more in *Bos indicus* than *Bos taurus* during winter (Randel, 2005), however, such effect of extreme cold has not been reported in buffaloes.

Lactation

High yielding cattle and buffalo shows significantly longer postpartum anestrus period (Harrison et al., 1989; El-Azab et al., 1984) or weaker signs of estrus (Harrison et al., 1990). El-Fadaly (1980) reported that buffalo producing >8 liters milk per day had longer postpartum anestrus (107±36 days) than those producing <8 liters per day (77±30 days). However, others have reported no significant correlation between milk yield and postpartum anestrus period (El-Keraby et al., 1981; Kawthar et al., 1985).

Suckling

Suckling suppress the postpartum ovarian activity both in cattle (Wiltbank and Cook, 1958; Oxenreider and Wagner, 1971; Stagg et al., 1998; Quintansa et al., 2009) and buffalo (El-Fadaly, 1980; Honnapagol et al., 1993), resulting into extended postpartum anestrus period. Moreover, postpartum anestrus is longer in continuously suckled than restricted or partial suckled cow and buffaloes (Bastidas et al., 1984; Nordin and Jainudeen, 1991). Postpartum anestrus is longer in suckled beef cow than milked dairy cow. Suckling stimulates prolactin, cortisol and oxytocin secretion that have negative effect upon GnRH-LH axis. Higher level of these hormones suppresses the GnRH secretion and increases the concentration of endogenous opioid peptides; β -endorphin (Malven et al., 1986) thus, ultimately reduces the LH pulse frequency (William, 1990) which delays resumption of postpartum cyclicity.

Parasitic Infestations

Heavy parasitism is one of the stressful conditions and is more common in growing than in adult cattle. It affects the future productive and reproductive efficiency in infested animals (Heath et al., 1997). Parasitic infection like fascioliasis, theleriosis schistosomiasis and trypanosomiasis infection in animals cause anemia and weight loss and ultimately results into anestrus. Recently, it has been found that *Neospora caninum* infection (Neosporosis) is widely prevalent among dairy herds and has significant association with anestrus (Bruhn et al., 2013).

Genotype

The resumption of postpartum cyclicity depends upon species as well as breeds. The postpartum anestrus period is shorter in milked dairy cows as compared to suckled beef cows but suckled dairy cows have longer postpartum anestrus period than beef cows (Short et al., 1990). The period of postpartum anoestrus is usually longer in buffalo than the cattle (Dobson and Kamonpatana, 1986; Jainudeen and Hafez, 1993). How genotype affect the resumption of postpartum cyclicity is not fully understood, however, it may be due to physiological differences among breeds and

species, difference in milk production and feed intake (Short et al., 1990).

Parity

A longer postpartum anestrus period have been reported in primiparous than pluriparous buffaloes (Ali and El-Sheikh, 1983; Barkawi, 1984). Moreover, as the parity increases, the postpartum anestrus period decreases (El-Sheikh and Mohamed; 1965; El-Wishy and El-Sawaf, 1971; Shah et al., 1989; Mahdy et al., 2001). However, others have reported there is no correlation between parity and postpartum anestrus period (El-Fouly et al., 1976; Borghese et al., 1993).

Periparturient Diseases

Periparturient Diseases such as abnormal calvings, metritis, mastitis and ketosis also influence onset of postpartum cyclicity (Fonseca et al., 1983; Opsomer et al., 2000). Delayed uterine involution also holdup resumption of ovarian activity. Postpartum uterine infection (clinical or sub clinical) suppress GnRH release and possibly LH secretion (Peter and Bosu, 1988; Peter et al., 1990; Mateus et al., 2002), probably due to inflammatory response (Sheldon and Dobson, 2004; Herath et al., 2006; Williams et al., 2007) and thus, ovarian activity remains suppressed in uterine infections.

DIAGNOSIS OF ANESTRUS

History

Based on the information viz., failure of displaying the overt signs of estrus by the animals after attaining puberty or 60–90 days post-partum; symptoms of estrus shown with cyclicity which subsequently ceased and revert in to anestrus. Such cases are diagnosed when presented for pregnancy diagnosis. Many times, owners complaint that they are not able to detect estrus or have not seen any signs of estrus in that particular animal since long.

Progesterone Estimation

True anestrus is usually characterized by a lack of ovarian progesterone production (Peter et al., 2009). Presence of basal level (0.5–1 ng/ml) of progesterone in the blood samples at an interval of 8–10 days further confirms the diagnosis. If the concentration of progesterone is more than 1ng/ml, it is suggestive of presence of corpus luteum and anestrus in such situation might be due to unobserved estrus/silent estrus/persistent corpus luteum.

Per Rectal Examination

Pregnancy can be a prominent cause of anestrus and therefore must be ruled out by careful examination of ovary and uterus when any animals present for gynecological examinations. On per rectal examination, ovaries are smooth, small and inactive with the absence of corpus luteum in true anestrus cattle and buffaloes (Agarwal et al., 2004), however, follicles may develop up to prematuration stage and get atretic (Roche et al., 1998; Ghuman et al., 2010). Functional corpus luteum can be palpated in case of silent estrus/unobserved as well as in anestrus due to persistent corpus luteum.

Ultrasonography

Ovarian pathology which is not accurately determined by per rectal palpation can be visualized by ultrasonography. Different stages of follicular growth and type of anestrus can easily be detected by ultrasonography. Transrectal ultrasonographic examination of anestrus buffaloes cows which are not seen in oestrus for 60 or more days postpartum at 12 days revealed 45% inactive ovaries (true

anestrus), 55% silent ovulation or missing heat (Rahman et al., 2012). It can also differentiate between persistent follicle and persistent CL.

TREATMENT

Anestrus can be treated according to their cause, however; there is no single panacea to correct it. Various therapeutic agents including hormonal and non-hormonal compounds have been used extensively for the restoration of cyclicity in anestrus cattle and buffalo by several workers with varying degree of success (Glotra et al., 1970; Deshpande et al., 2000, Agarwal et al., 2001, Kumar et al., 2005). In order to ensure effective treatment, the health and nutritional status of the animals must be in good conditions. Besides deworming, the supplementation of vitamins, minerals and antioxidants in feed are useful to improve health status of the animals.

NON HORMONAL TREATMENTS

Plant Based Heat Inducers

Plants have been used for the treatment of animals since long back. Plants synthesize varieties of phytochemicals such as alkaloids, glycosides, terpenes and tannins (secondary metabolites) as a part of their normal metabolic activity and many of these have therapeutic actions when consumed by animals. Many plants are rich source of vitamins and minerals whereas some have estrogenic property which is useful in restoration of cyclicity in anestrus animals. Almost all the parts of plant such as seeds, berries, roots, leaves, bark and flowers have been used as therapeutic agents either directly (crude drugs) or their active principles, after separation through various chemical process. Many plants such as *Murraya koenigii* (curry leaves), *Nigella sativa* (kalonji), *Abroma augusta* (Ulatkambal), *Saraca asoca* (Ashoka), *Trigonella foenum-graecum* (Methi), *Bambusa aruninacea*, *Carica papaya*, *Asparagus recemosus*, *Leptadenia reticulata*, *Courupita guianensis*, *Pergulacia daemia*, *Semecarpus anacardium cucumber*, and *jute plants* either alone or in combinations have been fed to treat the anestrus animals with variable response on induction of estrus (Kabir et al., 2001; Das et al., 2002; Mehrotra, 2002; Mishra et al., 2002; Rajkumar et al., 2008; Kumar and Punniamurthy, 2009). Kabir et al. (2001) reported 50% estrus induction in anestrus buffaloes using mixture of *Abroma augusta* (root) and *Nigella sativa* (seed) in 2:1 ratio. Rajkumar et al. (2008) reported higher success rate in anestrus cattle i.e. 83.33 and 66.66% using Methi seed (@ 200g/day/cow for 20 days) and bark of Ashoka tree (@ 50g/day/cow for 20 days), respectively.

Indigenous herbal preparations such as Prajana HS (Indian Herbs), Janova (Dabur), Sajani (Sarabhai), Heat up (Century) Heat raj (Ranjan), Fervivet (Ar Ex Labs) and Aloes compounds (Alarsar) are commercially available and effective in restoration cyclicity with good success rates (Deshpande and Sane, 1977; Patil et al., 1983; Dutta et al., 1988; Sudhir Chandra Reddy et al., 1990; Hussain et al., 2009). These formulations are potent combinations of herbs, formulated to induce ovarian activity.

Utero-Ovarian Massage

Utero-ovarian massage is the oldest, simplest, cheapest and effective method to induce estrus in anestrus cattle and buffaloes (Romaniuk, 1973; Zduńczyk et al., 1992; Rahawy, 2009; Mwaanga et al., 2010). Estrus induction in cattle and

buffalo varies between 40 to 80% following utero-ovarian massage daily/on alternate day/weekly for 3–4 weeks (Mwaanga et al., 2004; Naidu et al., 2009; Gupta et al., 2011). The mechanism by which ovarian massage induces cyclicity is not clearly understood, however, probable mechanism includes: activation of intrinsic intra-ovarian factors; enhancement of blood circulation to the ovaries and uterus that increases the availability of hormones and growth factors; stimulation of local oxytocin production by the ovaries which consequently influence local blood circulation and luteolysis, if CL is present (Romaniuk, 1973; Lobb and Dorrington, 1992; Monget and Monniaux, 1995; Mwaanga et al., 2010).

Lugol's Iodine

Lugol's iodine treatment is cheaper and effective means of management of anestrus but response has been variable (45 to 91.7%) among cattle and buffaloes (Porwal et al., 1976; Ryot et al., 1990; Agarwal and Pandit, 1991; Reddy et al., 1994; Megahed et al., 1995; Kendre and Bhosker, 1996; Singh and Thakur, 1999; Tapas et al., 2000; Tomar, 2004; Gupta et al., 2011). Lugol's iodine solution (5%) has traditionally been used as a cervical paint. It is presumed that painting of Lugol's iodine on posterior part of the cervix causes local irritation and brings about reflux stimulation at anterior pituitary for secretion of gonadotrophins and consequently cyclicity. Lugol's iodine is an irritating solution and intrauterine infusion of Lugol's solution (0.5 to 1.0%) causes hyperemia (enhanced circulation) of uterine mucosa resulting into degree of iodine absorption from uterus. The absorbed iodine probably increases the metabolic rate of body through stimulating the thyroid hormone secretion (Sanchez, 1995). Increased metabolic rate trigger the ovarian functions by enhancing the energy utilization (El-Shahat and Badr, 2011). Injectable Lugol's iodine has also been used with the same assumption (Sarkar, 2005). Another probable mechanism of intrauterine use of Lugol's iodine is that it acts as chemical curator (due to its irritating nature) and replaces the uterine mucosa with new tissue. The newly formed tissues of endometrium release luteolytic factors ($PG_{F_{2\alpha}}$) that reaches to the corpus luteum via utero-ovarian pathway and causes luteolysis (Gupta et al., 2011). Thus, it initiate the estrus cycle, if anestrus is due to PCL. Now-a-days, the use of Lugol's solution is not being recommended for treatment of anestrus due to its irritating nature and damaging effect on endometrium.

HORMONAL TREATMENTS

Estrogens Based Treatment

Exogenous administration of estrogen produce estrus signs in anestrus animals with or without concurrent ovulation. In presence of dominant follicle, estrogen administration results in expression of estrus and ovulation because of its positive feedback effect over pituitary for LH surge. For this reason, it has been used to induce ovulation (Saiduddin et al., 1968; Peters, 1984; Garcia-Winder et al., 1988; McDougall et al., 1994) and to reduce postpartum anestrus period. Conversely, estrogen induces anovulatory estrus in absence of dominant follicle. Estrogens have also been shown to cause luteolysis in ruminants (Wiltbank, 1966; Brunner et al., 1969) probably through stimulating the prostaglandin secretion from endometrium as well decreasing the level of circulating LH. One or two doses of intramuscular injections estradiol (3–10mg) or estrone (5–

15mg) at three days interval can be used to regresses the retained corpus luteum associated with pyometra, mummification and mucometra. However, it should be kept in mind that high and prolonged dose of estrogens leads to cystic ovaries, abnormal motility and peristalsis of oviduct that may results in introduction of infections to the ovarian bursa through oviduct, causing ovaritis and adhesions. These side effects of estrogens limit its use for induction of estrus in anestrus animals.

Progesterone Based Treatment

Exogenous administration of progesterone mimics the luteal phase of the estrus cycle by exerting negative feedback effect over hypothalamus and pituitary for LH release. Upon withdrawal of progesterone, the normal follicular phase of the cycle is stimulated. However, for such treatment seem to be effective, abrupt decrease in progesterone level is required at the end of treatment. Intravaginal progesterone releasing devices such as PRID (progesterone-releasing intravaginal device), CIDR (controlled internal drug release) and CueMate are effective in restoration of cyclicity in anestrus animals (Ramamohana Rao, 1981; Singh et al., 1988; Singh et al., 2010; Azawi et al., 2012). Upon withdrawal, the concentration of progesterone decline abruptly and onset of estrus and ovulation occurs within 2–8 days after the end of treatment (Agarwal and Tomar, 2003; Hafez and Hafez, 2008). Ear implants (Crestar and Synchronate-B) also produce required abrupt decrease in progesterone concentration at the end of treatment. Progesterone therapy alone is not particularly effective for the treatment of anovulatory anestrus; hence other hormones have been incorporated in most of the progesterone based therapy. To achieve better responses, the intravaginal devices or ear implants are generally used for 7 to 9 days, combined with other hormones (prostaglandins, GnRH, PMSG/eCG and estradiol) towards the end of progesterone treatment and estrus induction rate has been reported between 80 to 100% by most of the workers (Lakra et al., 2003; Rhodes et al., 2003; Kumar and Mandape, 2004; Nayak et al., 2009; Singh et al., 2010; Azawi et al., 2012).

Feeding of oral progestational compounds (10–14 days) such as MAP (6-methyl-17-acetoxypregesterone), CAP (6-chloro-6-dihydro-17-acetoxypregesterone), MGA (Melengestrol acetate) and DHPA (16 α -17-Dihydroxyprogesterone acetophenide) (Hansel and Malven, 1960; Hansel et al., 1966; Wiltbank et al., 1967; Shanker et al., 1988) and long term (4–14 days) intramuscular injections of progesterone @ 50–100 mg either alone or in combination with other hormones (Willet et al., 1950; Ulberg and Lindlay, 1960; Agarwal., 1983) have been used for induction/synchronization of estrus, however, both oral and parenteral routes are impractical for inducing the cyclicity as concentration of progesterone does not decline abruptly at the end of treatment.

The estrus response is better with progesterone based treatment, however, conception rates are usually low at induced estrus especially after long term progesterone therapy probably due to altered follicular growth, high rate of follicular atresia, poor sperm transport, failure of fertilization, low cleavage rate, poor transport of fertilized ova and early embryonic death (Jainudeen and Hafez, 1966; Hawk, 1971; Lamond et al., 1971; Wordinger et al., 1976).

Gonadotropic Releasing Hormone (GnRH) Based Treatment

The single intramuscular injection of GnRH analogue (10 to 20µg Buserelin) has been used effective in induction of estrus and concurrent ovulation with variable response (45.5 to 87.5%) within 4–22 days (Dhoble and Gupta, 1986; Pattabiraman et al., 1986; Sonwane et al., 1994; Narasimha Rao, 1997; Nautiyal et al., 1997; Markandeya and Patil, 2003; Prahalad et al., 2010). The variable response may be due to differential action of GnRH on different stages of follicular development. It induces ovulation, if mature follicle is present at the time of administration by inducing the LH surge. However, single injection of GnRH is not always effective in deep anestrus animals. On the contrary, it stimulate emergence of new follicular wave through enhanced secretion of FSH, thus effective in long term. Pulsatile/intermittent injections of small dose of GnRH (at every 2 hours, intravenously) has been tried in order to induce LH pulses (Edwards et al., 1983; Peters et al., 1985; Spicer et al., 1986), however, intermittent injection make this technique impractical.

To achieve better response, GnRH has been combined with other drugs such as phosphorus injection (Tonophosphan), prostaglandin, estradiol and progesterone (Shams et al., 1991; Rhodes et al., 2003; Sirmour et al., 2006). The Ovsynch protocol or GPG regimen (GnRH-PG-GnRH), developed by Pursley et al. (1995) to synchronize ovulations in dairy cows has been widely used to treat anestrus cattle and buffaloes and results are also promising. Under this protocol first injection of GnRH (at day 0) induces ovulation, if dominant follicle is present and if not luteinizes with emergence of new follicular wave 1 to 2 days later, PGF_{2α} injection given on day 7 regress the CL formed in response to first injection of GnRH and second injection GnRH on day 9 induces ovulation of new dominant follicle subsequently, all the treated animals are inseminated within 16–20 hours of second injection of GnRH.

Gonadotrophins Based Treatment

Pregnant mare serum gonadotrophin (PMSG) or equine chorionic gonadotrophin (eCG) is strong stimulator of ovarian activity because of its predominant FSH like activity. Therefore, it has been used extensively for superovulation. However, single intramuscular injection in low doses either alone or in combinations with others has been used successfully to treat anestrus cattle and buffaloes (Echternkamp, 1978; Wettemann, 1982; Dabas and Bardhan, 2006; Muneer et al., 2009; Prahalad et al., 2010; Kumar, 2012). PMSG prevent and reverse the process of atresia in small follicles (Moor et al., 1984), hence its use for management of anestrus in buffaloes in low doses could be used satisfactory as follicular atresia is very common in buffaloes. Human chorionic gonadotrophin (hCG) has also been used for management of anestrus with fair degree of success (Dabas and Bardhan, 2006).

Prostaglandin Based Treatment

Prostaglandin (PGF_{2α}) is the treatment of choice for persistent corpus luteum and sub estrus. Natural or synthetic analogue of PGF_{2α} as a single dose has been used with a reasonable degree of success for management of silent estrus in cattle and buffaloes (Nautiyal et al., 1998; Singh et al., 2001). It should be born in mind that PGF_{2α} is only effective between days 6–16 of the cycle and in the presence of active corpus luteum. An intramuscular

injection of 25mg (total dose) of natural PGF_{2α} or 250 to 500 micrograms of synthetic ones is required to regress the CL in both cattle and buffaloes. However, a lower dose of PGF_{2α} (5mg) are also effective to regress the CL through intra-vulvo-submucosal (IVSM) (Dhaliwal et al., 1988; Narasimha Rao and Venkatramaiah, 1990). Alternatively, Ovsynch protocol as describe by Pursley et al. (1995) may be used to treat sub estrus or unobserved estrus.

Insulin Based Treatment

Use of insulin for induction of estrus in animals either alone or in combination is a fairly recent development and results are very encouraging (Shukla et al., 2005a&b; Ramoun et al., 2007; Gupta et al., 2010). The recommended dose is 0.25 IU/kg body weight subcutaneously for 3–5 days. Use of GnRH or eCG pretreated with insulin has shown promising results for management of anestrus cattle (Shukla et al., 2005 a&b) and buffaloes (Gupta et al., 2010; Kumar et al., 2012; Ramoun et al., 2012). It has also been shown that insulin enhances the follicular growth in true anestrus buffalo which is prerequisite of GnRH to be effective (Ramoun et al., 2012). The treatment of true anestrus buffaloes under field condition was reported satisfactory when single intramuscular injection of PMSG (500 IU) was combined with subcutaneous injections of insulin @ 0.25IU/Kg body weight for five consecutive days (Kumar, 2012).

Anti-Prolactin Based Treatment

Hyper-prolactinaemia has been reported during summer in buffaloes that could be one of the reasons for summer anestrus in buffaloes. With this assumption anti-prolactin drug such as bromocriptine (Verma et al., 1992) has been tried. Melatonin is also known to suppress prolactin secretion (Wuliji et al., 2003). Moreover, melatonin has been reported as stimulator of both GnRH and gonadotrophin secretion in buffaloes. As the plasma concentration of melatonin is low during summer, Ghuman et al. (2010) has reported estrus induction and ovulation in all treated summer anestrus buffalo heifers using melatonin implants, however, time taken to induce estrus and ovulation was highly variable (4–36 days).

PREVENTION

Prevention of anestrus is preferable over treatment and can be achieved by maintaining the healthy status of the animals by adopting efficient farm managerial practices. Nutrition is probably most important factors, affecting ovarian activity. Special attention must be given to prevent the negative energy balance in high yielders. It can be achieved by providing adequate ration during pre- and postpartum period. The supplementation of vitamins, minerals and antioxidants in feed appeared to be promising in restoration of cyclicity (Baldi et al., 2000; Anita et al., 2003; Koley and Biswas, 2004; Selvaraju 2009; Sah and Nakao, 2010; Kumar, 2012).

As suckling decreases the LH pulsatility and prolongs the postpartum anestrus period, weaning could be one of the effective managerial tools to reduce postpartum anestrus period following restoration of pulsatile secretion of LH in postpartum animals. It has been shown that weaning (complete, temporary and partial) is associated with increased GnRH secretion, LH pulse frequency and more LH receptors on follicular cells followed by ovulation within few days. The level of opioid may also decrease due

to alleviation of physiological effects of suckling consequently, initiate normal ovarian cycles (Williams and Griffith, 1995) in cattle and buffaloes.

Biostimulation (induction of cyclicity through introduction of males into a group of females) activates LH secretion followed by ovulation (Gelez and Fabre-Nys, 2004) in females through olfactory and sensory cues. Exposure of postpartum cows (Tauck et al., 2010) and buffalo (Gokuldas et al., 2010) to a bull decreases the postpartum anestrus interval and advances the puberty in heifers. It can be elicited with vasectomized bulls, andronized females or steers. Though, it is a relatively inexpensive managemental tool, the response is inconsistent due to variability in parity, season, cow to bull ratio, body condition score at parturition and postpartum stages at which bulls are exposed. In cattle and buffaloes, the satisfactory response to biostimulation is achieved only when females are exposed to bulls immediately and continuously after parturition.

In addition to above preventive measures, certain managemental practices such as efficient detection of estrus either visually or by bull parading and other techniques, routine pregnancy diagnosis (40–60 days post breeding), prevention of occurrence of post-partum uterine infections and periparturient diseases (ketosis and mastitis), regular deworming and synchronization of estrus especially in buffaloes where heat detection is more difficult, could be advantageous in preventing the occurrence of anestrus.

CONCLUSION

Anestrus is a multi-causative factors associated problem affecting livestock enterprise to a great extent. Diagnosis of the condition needs to be prompt and at the earliest to prevent its occurrence for effective treatment. As such there is no single panacea to correct it. Further, research is needed especially at cellular and molecular level for better understanding of its etiology, diagnosis and therapeutics.

REFERENCES

Agarwal RG, RK Pandit (1991). Practical approach to therapeutic management of some commonly occurring reproductive disorders leading to infertility in bovines. *Vet.* 3(4): 219 – 225.

Agarwal SK, Tomer OS (2003). Reproductive technologies in buffalo. 2nd edn. A monograph published by communication centre, Indian Veterinary Research Institute, Izatnagar, India.

Agarwal SK, Patil RR, Purby LN (1983). Steroid induced ovarian cyclicity in anestrus cows. *J. Vet. Physiol. Allied Sci.* 2(1): 47 – 51.

Agarwal SK, Shanker U, Ansari MR (2004). Laboratory manual on Animal Gynaecology. Indian Veterinary Research Institute, Izatnagar, India.

Agarwal SK, Shanker U, Kumar S, Mohan G (2001). Ovarian cyclicity and progesterone profile in post-partum anoestrus cattle using synthetic progesterone, norgestomet regime. *Indian J. Anim. Sci.* 71: 1120 – 1123.

Agarwal SP, Buck NC (1968). Incidence of abnormal ovarian conditions and their impact on reproduction in breeding herd of cows. *Indian J. Vet. Sci.* 38: 533 – 540

Ali MA, El-Sheikh AS (1983). Postpartum ovulation in a herd of Egyptian buffaloes. *Indian J. Anim. Sci.* 53: 485 – 487.

Allrich RD (1994). Endocrine and neural control of estrus in dairy cows. *J. Dairy Sci.* 77: 2738 – 2744.

Anderson J (1936). Studies in reproduction in cattle. The periodicity and duration of oestrus. *Emp. J. Exp. Agric.* 4: 186.

Anita SN, Sangha SPS, Singh N (2003). Lipid peroxidation and antioxidant vitamins in postpartum anoestrus buffaloes supplemented with vitamin E and selenium. *Indian J. Dairy Sci.* 56: 33 – 37.

Awasthi MK, Kavani FS, Siddiquee GM, Sarvaiya NP, Derashri HJ (2007). Is slow follicular growth the cause of silent estrus in water buffaloes. *Anim. Reprod. Sci.* 99: 258 – 268.

Azawi OI, Ali MD, Oday SA, Salih A, Al-Hadad AS, Mouayad SJ, Abdul Hussien AS (2012). Comparative efficacy of different CIDR protocols

for the treatment of postpartum anoestrus in Iraqi buffaloes. *Vet. World.* 5(4): 201 – 205.

Badr HM (1993). Study of some reproductive aspects for postpartum period in Egyptian buffaloes. PhD dissertation, EI-Azhar University, Cairo, Egypt.

Baker LN, Woehling HL, Casida LE, Grummer RH (1953). Occurrence of estrus in sows following parturition. *J. Anim. Sci.* 12: 33.

Baldi A, Savoini G, Pinotti L, Monfardini E, Cheli F, Dell'Orto V (2000). Effects of vitamin E and different energy sources on vitamin E status, milk quality and reproduction in transition cows. *J. Vet. Med. Series A-Physiology Pathology Clinical Medicine.* 47(10): 599 – 608.

Barkawi AH (1984). Prepartum and postpartum hormonal pattern and reproductive activity in Egyptian buffaloes. Ph.D. Thesis. Cairo University.

Barkawi AH, Shafie MM, Mekawy Y, Aboul-Ela MB (1986). The use of serum and milk progesterone concentration to monitor post-partum ovarian activity in Egyptian buffaloes. *Buffalo. J.* 2 (2): 125 – 134.

Bartha R (1971). Studien zu Fragen der Zebu-Rinderzucht in den Tropen. Weltforum Verlag, Munich, Federal Republic of Germany. 172 pp.

Baruselli PS, Barnabe VH, Barnabe RC, Visintin JA, Molero-Filho JR, Porto R (2001). Effect of body condition score at calving on postpartum reproductive performance in buffalo. *Buffalo J.* 17 (1): 53 – 65.

Baruselli PS, Mucciolo RG, Visintin JA, Viana WG, Arruda RP, Madureira EH, Oliveira CA, Molero-Filho JR (1997). Ovarian follicular dynamics during the estrous cycle in buffalo (*Bubalus bubalis*). *Theriogenol.* 47: 1531 – 1547.

Bastidas P, Troconiz J, Verde O, Silva O (1984). Effect of restricted suckling on ovarian activity and uterine involution in Brahman cows. *Theriogenol.* 21: 525 – 532.

Bastos MR, Meschiatti MAP, Surjus RS, Prata AB, Lemes AP, Guardieiro MM, Santos FAP, Sartori R (2010). Estrus length and intensity in Bos taurus (holstein) vs. Bos indicus (Nelore) non lactating cows. *Reprod. Fert. Develop.* 23(1): 110 – 114.

Beam SW, Butler WR (1997). Energy balance and ovarian follicle development prior to the first ovulation postpartum in dairy cows receiving three levels of dietary fat. *Biol. Reprod.* 56: 133 – 142.

Bearden HJ, Fuquay JW, Willard ST (2004). *Applied Animal Reproduction*, Sixth Edition. Pearson Prentice Hall, Upper Saddle River, New Jersey, NY, U.S.A.

Beever DE, Hattan A, Reynolds CK, Cammell SB (2001). Nutrient supply to high yielding dairy cows. In: Diskin, M.G. (Ed.), *Fertility in the high producing dairy cow*. Occasional Publication No. 26, British Society of Animal Science, Edinburgh, pp. 119 – 131.

Bhagat S (1966). Artificial induction of lactation in freemartin. *Indian Vet. J.* 43: 218.

Bhalaru SS, Tiwana MS, Singh N (1987). Effect of body condition at calving on subsequent reproductive performance in buffaloes. *Indian J. Anim. Sci.* 57: 33 – 36.

Bharkad GP, Markandeya NM (2003). Incidence of bovine anoestrus. *Indian Vet. J.* 80: 190 – 191.

Bhattacharya P, Luktuke SN, Rao ASP, De SK (1954). Incidence of infertility under various causal groups in buffaloes cow in India. *Curr. Sci.* 23: 335.

Bhattacharyya, Buchoo (2008). Incidence of Reproductive Disorders of Cattle in Kashmir Valley. *Indian J. Dairy Sci.* 61: 165.

Bo GA, Baruselli PS, Martinez MF (2003). Pattern and manipulation of follicular development in Bos indicus cattle. *Anim. Reprod. Sci.* 78: 307 – 326.

Bonia KK (1981). A clinical note on the ovarian conditions of the rural cattle of Assam. *Indian J. Anim. Reprod.* 1: 17.

Borghese A, Barile VL, Terzano GM, Annicchiarico G, Debenedetti A, Malfatti A (1993). Anestrus length in Italian buffalo cows. In: *Prospects of Buffalo Production in the Mediterranean and Middle East*, Note I. Cairo, Egypt, pp 389 – 392.

Bruhn FRP, Daher DO, Lopes E, Barbieri JM, Rocha CMBM, Guimarães AM (2013). Factors associated with seroprevalence of Neospora caninum in dairy cattle in southeastern Brazil. *Trop. Anim. Health Prod.* 45(5): 1093 – 1098.

Brunner MA, Donaldson LE, Hansel Wm (1969). Exogenous hormones and luteal function in hysterectomized and intact heifer. *J. Dairy sci.* 52(11): 1849.

Butler WR, Smith RD (1989). Interrelationships between energy balance and postpartum reproductive function in dairy cattle. *J. Dairy Sci.* 72: 767 – 783.

Campbell CS, Wood R, Kelly M (1999). *Social Capital and Health*. Health Education Authority, London, UK.

Canfield RW, Butler WR (1990). Energy balance and pulsatile LH secretion in early postpartum dairy cattle. *Domest. Anim. Endocrinol.* 7: 323–330.

Chauhan FS, Takkar OP, Tiwana MS (1976). A note on incidence of gestational estrus in buffaloes and cattle. *Indian Vet. J.* 53: 77.

- Connor HC, Houghton PL, Lemenager RP, Malven PV, Parfet JR, Moss GE (1990). Effect of dietary energy, body condition and calf removal on pituitary gonadotropins, gonadotropin-releasing hormone (GnRH) and hypothalamic opioids in beef cows. *Domest. Anim. Endocrinol.* 7: 403 – 411.
- Cook DL, Smith CA, Parfet JR, Youngquist RS, Brown EM, Garverick HA (1990). Fate and turnover rate of ovarian follicular cysts in dairy cattle. *J. Reprod. Fert.* 90: 37 – 46.
- Crowe MA, Padmanabhan V, Mihm M, Beitins IZ, Roche JF (1998). Resumption of follicular waves in beef cows is not associated with periparturient changes in follicle-stimulating hormone heterogeneity despite major changes in steroid and luteinizing hormone concentrations. *Biol. Reprod.* 58: 1445 – 1450.
- Dabas YPS, D Bardhan (2006). Oestrus and fertility after gonadotrophic treatment in anestrus buffaloes. *Int. Polivet.* 4(2): 162 – 165.
- Damodaran S (1956). Studies on genital pathology of the female buffaloes M.Sc. Thesis Madras University, Madras.
- Danell B (1987). Oestrus behaviour, ovarian morphology and cyclical variation in the follicular system and endocrine pattern in Water buffalo heifers. Tese de doutorado, Universidade de Uppsala, Sweden.
- Das P, Das SK, Arya HP, Subba Reddy G, Mishra A (2002). *Veterinary Science and Animal Husbandary. In: Inventory of indigenous technical knowledge in agriculture, Document1, ICAR, New Delhi.* Pp. 185 – 285.
- De Alba JE, Corta V, Ulloa G (1961). Influence of natural service on length of oestrus in the cow. *Anim. Prod.* 3: 327.
- Deshpande BR, Sane CR (1977). Prolonged postpartum anestrus condition in cattle. *Indian Vet. J.* 54: 63 – 68.
- Deshpande RS, Dhoble RL, Sawale AG (2000). Efficacy of indigenous drugs in the treatment of post-partum anoestrus in buffaloes. *Indian J. Anim. Reprod.* 21: 115 – 116.
- Dessouky F, Juma KH (1973). Infertility problems among cows and buffalo in Iraq. *Indian J. Anim. Sci.* 43: 187 – 192.
- Dhaliwal GS, Sharma RD, Singh G (1988). Efficacy of prostaglandin F₂-alpha administration for inducing estrus in buffalo. 29(6): 1401 – 1406.
- Dhoble RL, Gupta SK (1986). Response to synthetic gonadotrophin (GnRH) in anestrus cow. *Theriogenol.* 25: 759.
- Diskin MG (2008). Heat Watch: a telemetric system for heat detection in cattle. *Vet. Q.* 30: 37 – 48.
- Diskin MG, Mackey DR, Roche J, Sreenan JM (2003). Effects of nutrition and metabolic status on circulating hormones and ovarian follicle development in cattle. *Anim. Reprod. Sci.* 78: 345 – 370.
- Dobson H, Kamopatana M (1986). A review of female cattle reproduction with special reference to a comparison between buffaloes, cows and zebu. *J. Reprod. Fert.* 77: 1 – 36.
- Dutta JC, Deka KC, Rahman S (1988). Treatment of anoestrus cattle with aloes compound. *Indian J. Anim. Health.* 27(2): 171 – 172.
- Dziuk PJ, Bellows RA (1983). Management of reproduction of beef, sheep and pigs. *J. Anita. Sci.* 57 (Suppl. 2): 355.
- Easdon MP, Chesworth JM, Aboul-Ela MB, Henderson GD (1985). The effect of undernutrition of beef cows on blood hormone and metabolite concentrations post-partum. *Reprod. Nutr. Dev.* 25: 113 – 126.
- Echternkamp SE (1978). Stimulation of estrogen and luteinizing hormone secretion in postpartum beef cows. *J. Anim. Sci.* 47: 521 – 531.
- Edwards S, Roche JF, Niswender GD (1983). Response of suckling beef cows to multiple, low-dose injections of GnRH with or without progesterone pretreatment. *J. Reprod. Fert.* 69: 65 – 72.
- El-Azab EA, Mansour H, Heshmat H, Shawki G (1984). The postpartum period and the future fertility of the Egyptian buffalo. In: *Proceedings of the 10th International Congress on Animal Reproduction and AI, vol. III, No. 424, Urbana, p. 3*
- El-Fadaly MA (1980). Effect of suckling and milking on the breeding efficiency of buffaloes. I. First postpartum estrus. *Vet. Med. J.* 28: 399 – 404.
- El-Fouly MA, Kotby EA, El-Sobhy HE (1976). Postpartum ovarian activity in suckled and milked buffaloes. *Theriogenol.* 5: 69 – 81.
- El-Keraby F, Aboul-Ela MB, Bedeir LH (1981). The effect of diet on postpartum reproductive traits in buffaloes. *Agric. Res. Rev.* 59: 1 – 12.
- El-Sawaf SA, Shalaby AS, Kamal TH (1979). The effect of heat stress on the performance of Friesian heifers. *Zagazig Vet. J.* 2: 185 – 192.
- El-Shahat KH, Badr A (2011). Comparative Study on Efficacy of Different Medicaments on Postpartum Anestrus Dairy Cows. *J. Appl. Biol. Sci.* 5(3): 59 – 63.
- El-Sheikh AS, Mohamed AA (1965). The reproductive performance of the Egyptian buffalo. *Egypt. J. Anim. Prod.* 5: 99 – 117.
- El-Wishy AB (2007). The postpartum buffalo II. Acyclicity and anestrus. *Anim. Reprod. Sci.* 97: 216 – 236.
- Ferguson JD (2001). Nutrition and reproduction in dairy herds. In: *Proc. Intermountain Nutr. Conf. Utah State Univ. Logan.* p. 65
- Fonseca FA, Britt JH, McDaniel BT, Wilk JC, Rakes AH (1983). Reproductive traits of Holsteins and Jerseys. Effects of age, milk yield, and clinical abnormalities on involution of cervix and uterus, ovulation, estrous cycles, detection of estrus, conception rate, and days open. *J. Dairy Sci.* 66: 1128 – 1147.
- Forde N, Beltman ME, Lonergan P, Diskin M, Roche JF, Crowe MA (2011). Oestrous cycles in *Bos taurus* cattle. *Anim. Reprod. Sci.* 124: 163 – 169.
- Fortune JE (2003). The early stage of follicular development: activation of primordial follicles and growth of preantral follicles. *Anim. Reprod. Sci.* 78: 135 – 163.
- Fortune JE, Rivera GM, Yang MY (2004). Follicular development: the role of the follicular microenvironment in selection of the dominant follicle. *Anim. Reprod. Sci.* 82 – 83: 109 – 126.
- Garcia-Winder M, Lewis PE, Inskip EK (1988). Ovulation in postpartum beef cows treated with estradiol. *J. Anim. Sci.* 66: 1 – 4.
- Garverick HA (1997). Ovarian follicular cysts in dairy cows. *J. Dairy Sci.* 80: 995 – 1004.
- Garverick HA (2007). Ovarian follicular cysts. In: *Current Therapy in Large Animal Theriogenology* (eds R. S. Youngquist and W. R. Threlfall). Saunders Elsevier, St. Louis, MO. p. 379 – 383.
- Gelez H, Fabre-Nys C (2004). The “male effect” in sheep and goats: a review of the respective roles of the two olfactory systems. *Horm. Behav.* 46: 257 – 271.
- Ghuman SP, Singh J, Honparkhe M, Dadarwal D, Dhaliwal GS, Jain AK (2010). Induction of ovulation of ovulatory size non-ovulatory follicles and initiation of ovarian cyclicity in summer anoestrus buffalo heifers (*Bubalus bubalis*) using melatonin implants. *Reprod. Domest. Anim.* 45(4): 600–607.
- Ghuman SPS, Singh J, Honparkhe M, Dadarwal D, Dhaliwal GS, Singh ST (2010). Fate of dominant follicle in summer anoestrus buffaloes. *Indian J. Anim. Reprod.* 31(2): 7 – 10.
- Gilbert RO, Bosu WTK, Peter AT (1990). The effect of *Escherichia coli* endotoxin on luteal function in Holstein heifers. *Theriogenol.* 33: 645 – 651.
- Ginther OJ, Kot K, Kulick LJ, Martin S, Wiltbank MC (1996). Relationships between FSH and ovarian follicular waves during the last six months of pregnancy in cattle. *J. Reprod. Fert.* 108: 271 – 279.
- Glotra AP, Tyagi RPS, Singh RP (1970). Effect of double application of Lugol's solution and oral feeding of potassium iodide on induction of oestrus, ovulation and conception in buffaloes and heifers. *Indian J. Anim. Sci.* 40: 437.
- Gokuldas PP, Yadav MC, Kumar H, Singh G, Mahmood S, Tomar AKS (2010). Resumption of ovarian cyclicity and fertility response in bull-exposed postpartum buffaloes. *Anim. Reprod. Sci.* 121: 236 – 240.
- Grimard B, Humblot P, Ponter AA, Mailot JP, Sauvart D, Thibier M (1995). Influence of postpartum energy restriction on energy status, plasma LH and estradiol secretion and follicular development in suckled beef cows. *J. Reprod. Fert.* 104: 173 – 179.
- Gupta M, Verma HK, Kasrija R (2010). Treatment of anoestrus in jersey crossbred heifers. *Indian J. Field Vet.* 6 (2): 19 – 20.
- Gupta R, Thakur MS, Sharma A (2011). Estrus induction and Fertility response in true anoestrus buffaloes using lugol's iodine. *Vet. World.* 4(2): 77 – 78.
- Gupta V, Thakur MS, Agrawal RG, Quadri MA, Shukla SN (2010). Effect of pretreatment with Insulin on ovarian and fertility response in true anoestrus buffaloes to Gonadotrophin-Releasing hormone. *Buffalo Bull.* 29(3): 172 – 179.
- Hafez B, Hafez ESE (2008). *Reproduction in farm animals*, 7th ed. Blackwell Publishing Limited. p 407.
- Hamilton SA, Garverick HA, Keisler DH, Xu ZZ, Loos K, Youngquist RS, Salfen BE (1995). Characterization of ovarian follicular cysts and associated endocrine profiles in dairy cows. *Biol. Reprod.* 53: 890 – 898.
- Hansel W, Malven PV (1960). Estrus cycle regulation in beef cattle by orally active progestational agent. *J. Anim. Sci.* 19(4): 1324.
- Hansel W, Donaldson LE, Wagner WC, Brunner MA (1966). A comparison of estrous cycle synchronization methods in beef cattle under feedlot conditions. *J. Anim. Sci.* 25: 497.
- Harrison RO, Ford SP, Young JW, Conley AJ, Freeman AE (1990). Increased milk production versus reproductive and energy status of high producing dairy cows. *J. Dairy Sci.* 73: 2749 – 2758.
- Harrison RO, Young JW, Freeman AE, Ford SP (1989). Effects of lactational level on reactivation of ovarian function, and interval from parturition to first visual estrus and conception in high-producing Holstein cows. *Anim. Prod.* 49: 23 – 28.
- Hawk HW (1971). Sperm destruction in the sheep vagina. *J. Anim. Sci.* 33: 255 (Abstr.).
- Heath HL, Blagburn BL, Elsasser TT, Pugh DG, Sanders LG, Sartin E, Steele B, Sartin JL (1997). Hormonal modulation of the physiologic responses of calves infected with *Eimeria bovis*. *Am. J. Vet. Res.* 58: 89 – 896.
- Hegazi MA, El-Wishy AB, Youssef AH, Awadalla SA, Teleb HM (1994). Interrelationship between pre- and/or post-partum feeding levels, blood constituents and reproductive performance of buffaloes. In: *Proceedings of the Fourth World Buffalo Congress, vol. III, San Paulo, Brazil, pp. 632 – 633.*

- Herath S, Dobson H, Bryant CE, Sheldon IM (2006). Use of the cow as a large animal model of uterine infection and immunity. *J. Reprod. Immunol.* 69: 13 – 22.
- Hidiroglou M (1979). Manganese in ruminant nutrition. *Can. J. Anim. Sci.* 59: 217.
- Honnapagol SS, Muregeppa A, Biradar US, Mallikarjunappa S (1993). Postpartum reproductive performance in suckled and non-suckled Surti buffaloes. *Indian Vet. J.* 70: 470 – 471.
- Hussain J, Bhat AS, Shaheen M, Islam R (2009). Comparative response of vitamin-mineral vs herbal therapy in alleviating post-partum true anestrus in dairy cows. *Indian J. Anim. Res.* 43 (3): 222 – 223.
- Hussain J, Bhat AS, Shaheen M, Islam R (2009). Comparative response of vitamin-mineral vs herbal therapy in alleviating post-partum true anestrus in dairy cows. *Indian J. Anim. Res.* 43(3): 222 – 223.
- Hussain PM (1984). Studies on the incidence of reproductive disorders of bovines in a part of southern Karnataka. *Livestock Advisor.* 9: 13.
- Iyer CPN, Nair KP, Sudarsan V, Madhvan E, Mothai E, Nair MS, Vijay KV, Joseph M (1992). Reproductive disorders of crossbred cows in Kerala. *Indian J. Anim. Reprod.* 13: 64 – 68.
- Jainudeen MR, Hafez ESE (1966). Control of estrus and ovulation in cattle with orally active progestin and gonadotrophin. *J. Fert. II(1)*: 47.
- Jainudeen MR, Hafez ESE (1993). Cattle and buffalo. In: Hafez, E.S.E. (Ed.), *Reproduction in Farm Animals*, 6th ed. Lea and Febiger, Philadelphia, USA. p 315 – 329.
- Kabir KK, Rawal CVS, Ansari MR, Varshney JP, Srivastava RS (2001). Comparative efficacy of herbal preparations in the management of anoestrus in non-descript rural buffaloes. *Indian J. Anim. Reprod.* 22(2): 143 – 145.
- Kaikini AS (1974). Studies on bovine gynaecology-gonads and reproductive tract of Berari buffaloes. Ph.D. Thesis, Punjab Agric. Univ.
- Kaikini FS, Fasihuddin M (1984). Incidence of gestational oestrus I Sahiwal and Gir cow. *Indian J. Anim. Sci.* 54 (9): 886 – 887.
- Kawate N (2004). Studies on the regulation of expression of luteinizing hormone receptor in the ovary and the mechanism of follicular cyst formation in ruminants. *J. Reprod. Dev.* 50: 1 – 8.
- Kawthar AM, Youssif RH, Khattab AS (1985). Individual factors affecting the interval between calving and first service in Egyptian buffaloes. *Egypt. J. Anim. Prod.* 25: 319 – 327.
- Kezele P, Nilson EE, Skinner MK (2002). Insulin but not insulin like growth factors-1 promotes the primordial to primary follicle transition. *Mol. Cell. Endocrinol.* 198 (1 – 2): 37 – 43.
- Khan BU, Luktuke SN (1967). Studies on incidence of anoestrus and repeat breeding in Harian cows. *Indian Vet. J.* 44: 678 – 683.
- King GJ, Hurnik JF, Robertson HA (1976). Ovarian function and estrus activity in dairy cows during early lactation. *J. Dairy Sci.* 42: 688 – 92.
- Knickerbocker JJ, Thatcher WW, Foster DB, Wolfenson D, Bartol FF, Caton D (1986). Uterine prostaglandin and blood flow responses to estradiol-17b in cyclic cattle. *Prostaglandins.* 31: 757 – 776.
- Kodagali SB (1968). A note on reproductive disorders of farm animals in Saurashtra. *Indian J. Vet. Sci. Anim. Husband.* 38: 286 – 287.
- Kodagali SB (1969). Bilateral ovarian hypoplasia in Gir heifer. *Indian Vet. J.* 46: 289 – 290.
- Kodagali SB (1974). Report on study of infertility in cattle (1965 – 1969), Anand, Gujarat, India. *Vet. Bull.* 45: 1285.
- Kodagali SB (1974). Reports on study of infertility in cattle (1965 – 1969) Anand, Gujarat. *Indian Vet. Bull.* 45: 1285.
- Kodagali SB, Kerur VK (1968). Genital abnormalities in Jaffarabadi buffalo cow Gujarat Vet. J. 2: 29.
- Koley S, Biswas P (2004). Effect of mineral supplementation on the performance of anoestrous cows. *Indian J. Anim. Nutr.* 21(4): 268 – 270.
- Kulkarni MD, Khanvilkar AV, Bansod RS (2002). Incidence of reproductive disorders in indigenous and crossbreds in an organized farm. *Indian Vet. J.* 79: 1196 – 97.
- Kumar PR (2012). Incidence of anoestrus in buffaloes and its therapeutic management using PMSG and Insulin. M.V.Sc. Thesis, MPCCVV, Jabalpur.
- Kumar H, Mandape MK (2004). Fertility management in rural buffaloes by hormonal therapies during the summer season. *Buffalo Bull.* 23(2): 30 – 33.
- Kumar PR, Shukla SN, Purkayastha RD (2013). Economical analysis of the estimated cost of management of anoestrus buffaloes under field conditions using different hormonal and non-hormonal strategies. *J. Anim. Health Prod.* 1(4): 39 – 41.
- Kumar PR, Shukla SN, Shrivastava OP, Purkayastha RD (2013). Incidence of postpartum anoestrus among buffaloes in and around Jabalpur. *Vet. World.* 6(9): 716 – 719.
- Kumar S, Agarwal SK (1986). Studies on ovarian dysfunction in rural bovines. *Indian Vet. Med. J.* 10: 11 – 22.
- Kumar S, Punniamurthy N (2009). Estrus induction by supplementation of *Murraya koenigii* in anoestrus heifers. *Indian J. Anim. Reprod.* 30(2): 66 – 67.
- Kumar S, Misra AK, Singh M (2005). Induction of oestrus in post-partum anoestrous cows with Creastar, GnRH and hCG. *Indian J. Anim. Sci.* 75: 22 – 24.
- Kutty I, Ramachandran K. (2003). Bovine infertility: a field oriented categorisation based on investigation among crossbred cattle in a district of Kerala. *Indian J. Anim. Sci.* 73: 155 – 157.
- Lagerlof N (1951). Hereditary forms of sterility in Swedish cattle breed. *Fert. Steril.* 2(3): 230.
- Lagerlof N, Boyd H (1953). Ovarian hypoplasia and other abnormal conditions in the sexual organs of cattle of Swedish Highland breed: Results of postmortem examination of over 6,000 cows. *Cornell Vet.* 43: 52.
- Lakra BS, Luthra RA, Khar SK, Nanda T, Baniwal BS (2003). Induction of cyclicity in anoestrous buffaloes during non-breeding season. *Intas Polivet* 4(2): 162 – 166.
- Lalman DL, Keisler DH, Williams JE, Scholljegerdes EJ and Mallet DM (1997). Influence of postpartum weight and body condition change on duration of anoestrus by undernourished suckled beef heifers. *J. Anim. Sci.* 75: 2003 – 2008.
- Lamond DR, Dickey JF, Henricks DM, Hill JR, Leland TM (1971). Effect of progestin on the bovine ovary. *J. Anim. Sci.* 33: 77.
- Lobb DK, Dorrington J (1992). Intraovarian regulation of follicular development. *Anim. Reprod. Sci.* 28: 343 – 354.
- Lopez H, Satter LD, Wiltbank MC (2004). Relation between level of milk production and estrus behavior of lactating dairy cow. *Anim. Reprod. Sci.* 81: 209 – 223.
- Lucy MC (2001). Reproductive Loss in High-Producing Dairy Cattle. *J. Dairy Sci.* 84: 1277 – 1293.
- Lucy MC (2003). Mechanisms linking nutrition and reproduction in post-partum cows. *Reprod. Suppl.* 61: 415 – 427.
- Lucy MC, Bilby CR, Kirby CJ, Yuan W, Boyd CK (1999). Role of growth hormone in development and maintenance of follicles and corpora lutea. *J. Reprod. Fert. Suppl.* 54: 49 – 59.
- Luktuke SN, Arora RL (1967). Studies on cystic degeneration of ovaries in Haryana cows. *Ind. Vet.* 44: 947.
- Luktuke SN, Sharma C (1978). Studies on the incidence of true anoestrus in rural cattle and buffaloes. *Indian Vet. J.* 55: 940 – 942.
- Luktuke SN, Bhattacharyya AR, Singh SK, Khan BV (1973). Studies on aberration in functional activity of the ovaries in buffaloes. *Indian Vet. J.* 50: 876 – 881.
- Luktuke SN, Roy JD, Joshi SR (1964). Studies on gestational estrus in bovines. *Indian J. Vet. Sci.* 34: 41.
- Lyimo ZC, Nielen M, Ouweltjes W, Kruij TA, Van Eerdenburg FJ (2000). Relationship among estradiol, cortisol and intensity of estrous behavior in dairy cattle. *Theriogenol.* 53: 1783 – 1795.
- Lynn JE, McNutt SH, Casida LE (1966). Effects of intrauterine bacterial inoculation and suckling on the bovine corpus luteum and uterus. *Am. J. Vet. Res.* 27: 1521 – 1526.
- Mahdy AE, El-Shafie OM, Rigalaty HA (2001). Relative importance of some factors affecting performance traits in a herd of Egyptian buffaloes. *Alex. J. Agric. Res.* 46: 1 – 18.
- Malik PS, Sengar OPS, Singh SN (1960). Structure of abnormalities of female genitalia of Indian buffalo. *Agra Univ. J. Res.* 9: 271.
- Malven PV, Parfet JR, Gregg DW, Allrich RD, Moss GE (1986). Relationships among concentrations of opioid neuropeptides and luteinizing hormone in neural tissues of beef cows following early weaning. *J. Anim. Sci.* 62: 723 – 733.
- Markandeya NM, Patil AD (2003). Studies on hormonal therapies for induction of post-partum estrus in buffaloes. *Intas Polivet.* 4(11): 167 – 169.
- Markusfeld O (1987). Inactive ovaries in high-yielding dairy cows before service: aetiology and effect on conception. *Vet. Rec.* 121: 149 – 53.
- Markusfeld O, Galon N, Ezra E (1997). Body condition score, health, yield and fertility in dairy cows. *Vet. Rec.* 141: 67 – 72.
- Mateus L, da Costa LL, Bernardo F, Silva JR (2002). Influence of puerperal uterine infection on uterine involution and postpartum ovarian activity in dairy cows. *Reprod. Domest. Anim.* 37: 31 – 35.
- McCracken JA, Custer EE, Lamsa JC (1999). Luteolysis: a neuroendocrine-mediated event. *Physiol. Rev.* 79: 263 – 323.
- McDougall S, Macmillan KL, Williamson NB (1994). The effect of oestradiol-178 on the rising and plateau dominant follicle in anoestrous cows. *Theriogenol.* 41: 252 abstr.
- McLaughlin EA, McIver SC (2009). Awakening the oocyte: controlling primordial follicle development. *Reprod.* 137: 1 – 11.
- Megahed GA, Anwar MM, Salwa MS (1995). Thyroid activity in Egyptian buffaloes cows with ovarian inactivity and trails for treatment by intra-uterine infusion of lugol's iodine solution. In: 3rd scientific congress Egyptian society for cattle disease, 3–5 December, Assiut-Egypt., 2: 211 – 217.

- Mehrotra S (2002). Studies on ovarian function using certain medicinal plants in rats, goats and cattle. Ph.D. Thesis submitted to I.V.R.I. Izatnagar, Bareilly.
- Mishra BP, Nayak N, Mishra D (2002). Indigenous methods for animal health care. International seminar on traditional knowledge, health and environment, held during 23–24 February 2002, at Bubaneswar, Abs, pp. 124.
- Mittal D, Garg UK, Jatav GP, Shukla S, Sharda R (2010). Prevalence of different pathological affections of ovaries in buffaloes (*Bubalus bubalis*) in malwa region of madhya Pradesh. *Buffalo Bulletin*. 29(1): 39 – 42.
- Modi LC, Patel PA, Patel SP, Patel GG, Joshi AH, Suthar DN (2011). Prevalence of reproductive problems in buffalo in Mehsana Milk–Shed area of Gujarat. *Int. J. Ag. Vet. Med. Sci.* 5(4): 424 – 428.
- Monget P, Monniaux D (1995). Growth factors and control of folliculogenesis. *J. Reprod. Fert.* 49: 321 – 333.
- Montiel F, Ahuja C (2005). Body condition and suckling as factors influencing the duration of postpartum anestrus in cattle: A review. *Anim. Reprod. Sci.* 85: 1 – 26.
- Moor RM, Kruip TAM, Green D (1984). Intra-ovarian control of folliculogenesis: Limits to superovulation. *Theriogenol.* 21: 103.
- Muneer S, Sadasiva RK, Solmon RKG (2009). Efficacy of GnRH-PGF_{2α}, PMSG and PMSG + hCG in postpartum anestrus crossbred cows. *Indian J. Anim. Reprod.* 30(1): 7 – 9.
- Mwaanga AA, Zdunczyk S, Janowski T (2004). Comparative study on the efficacy of hormonal and non-hormonal treatment methods in ovarian dysfunction affected dairy cow. *Bull. Vet. Inst. Pull.* 48: 265 – 267.
- Mwaanga ES, Janowski T, Zdunczyk S, Simukoko H (2010). Ovarian massage: a simple, but useful tool to manage ovarian acyclicity in dairy cows. *Bull. Anim. Health Prod. Afr.* 58: 294 – 295.
- Naidu GV, Srinivas M, Krishna HNVV, Prasad DV (2009). Management of delayed puberty in graded murrha heifers under field conditions – a practical approach. *Buffalo Bull.* 28(4): 204 – 206.
- Naidu KV, Rao AR (1981). Incidence of infertility among crossbred cattle of Andhra Pradesh. *Indian J. Anim. Sci.* 51: 829 – 831.
- Naidu KV, Rao AR. (1981). Incidence of infertility among crossbred cattle in Andhra Pradesh. *Indian J. Anim. Sci.* 51: 829 – 831.
- Nair KP, Raja CKSV (1974). Studies on the pathological conditions in the reproductive organs of cows. I. Pathology of ovaries. *Kerala J. Vet. Sci.* 5: 82.
- Nanda AS, Brar PS, Prabhakar S (2003). Enhancing reproductive performance in dairy buffalo: major constraints and achievements. *Reprod.* 61(Suppl.): 27 – 36.
- Nanda AS, Ward WR, Dobson H (1991). Lack of LH response to oestradiol treatment in cows with cystic ovarian disease and effect of progesterone treatment or manual rupture. *Res. Vet. Sci.* 51: 180 – 184.
- Narasimha Rao AV, and Murthy AK (1972). Studies on reproductive disorders in cow in Andhra Pradesh *Indian Vet. J.* 53: 156.
- Narasimha Rao AV, Sreemannarayana O (1982). Clinical analysis of reproductive failure among female buffaloes under village management in Andhra Pradesh. *Theriogenol.* 18: 403.
- Narasimha Rao AV, Murthy ST (1980). Incidence of freemartinism in the Indian water buffalo. *Indian Vet. J.* 57: 783.
- Narasimha Rao AV, Venkatramiah P (1990). The effectiveness of smaller doses of prostaglandin F₂ alpha in increasing the reproductive efficiency on Ongole cattle. *Indian vet. J.* 67: 528.
- Narladkar BW, Bakshi SA, Pargaonkar DR, Dgraskar SU (1994). Incidence of various reproductive disorders in Deoni cows and their crossbreds. *Liv. Adv.* 19(5): 28 – 30.
- Narsimha Rao AV (1997). Treatment of anoestrus cows and buffaloes with gonadotropin releasing hormone used singly or in combination with oestradiol 17-β. *Indian. Vet. J.* 74: 938 – 939.
- Nautiyal H, Shanker U, Agarwal SK (1997). Effect of gonadotropin releasing hormone (GnRH) on induction of ovarian cyclicity in pubertal anoestrus buffalo heifers. *Indian J. Anim. Reprod.* 18: 13 – 14.
- Nautiyal H, Shanker U, Agarwal SK (1998). Synchronization of oestrus using double injection regimen of PGF_{2α} in buffaloes. *Indian Vet. Med. J.* 22: 99 – 100.
- Nayak HB, Mohanty BN (1986). Incidence of anestrus in crossbred cattle. *Indian J. Dairy Sci.* 39: 314 – 315.
- Nayak V, Agrawal RG, Shrivastava OP, Thakur MS (2009). Induction of estrus in true anoestrus buffaloes using Crestar implant alone and in combination with PMSG. *Buffalo Bull.* 28(2): 51 – 54.
- Noakes DE, Parkinson TJ, England GCW (2009). *Veterinary Reproduction and Obstetrics*, 9th ed. WB Saunders Company, London.
- Noakes DE, Wallace LM, Smith GR (1990). Pyometra in a Friesian heifer: bacteriological and endometrial changes. *Vet. Rec.* 126: 509.
- Nordin Y, Jainudeen MR (1991). Effect of suckling frequency on postpartum reproductive performance of swamp buffaloes. In: *Proceedings of the Third World Buffalo Congress*, vol. II, Sofia, Bulgaria, pp. 737 – 743.
- Opsomer G, Grohn YT, Hertl J, Coryn M, Deluyker H, de Kruij A (2000). Risk factors for post-partum ovarian dysfunction in high producing dairy cows in Belgium: a field study. *Theriogenol.* 53: 841 – 57.
- Oxenreider SL, Wagner WC (1971). Effect of lactation and energy intake on postpartum ovarian activity in the cow. *J. Anim. Sci.* 33: 1026 – 1031.
- Pandit RK (2004). Incidence of different kinds of reproductive disorders in livestock. *Indian J. Anim. Reprod.* 25: 35 – 36.
- Pargaonkar DR (1978). Studies on bovine gynaecology, fertility and problems of reproduction in native cross-bred cattle, Ph.D. Thesis, PKV, Akola.
- Pargaonkar DR, Bakshi SA (1987). Reproductive disorders of red kandhari cows and their crossbreds. *Indian J. Anim. Reprod.*, 8 (1): 18 – 19.
- Patel PM, Dhama AJ, Sarvaiya NP, Kavani FS (2007). Monitoring postpartum period in holstein friesian cows through plasma progesterone profile with and without hormone therapy. *Indian. J. Anim. Sci.* 77(1): 3 - 9.
- Patil JS, Bangalia, Sinha NS, Sharma AK and Chopra SC (1983). Clinical efficacy of prajana, lugol's paint and uteroovarian massage in anoestrus cows. *Indian Vet. J.* 60: 1019 – 1020.
- Pawshe CH, Ingawale MV, Deshmukh SG, Munde VK, Pawshe MD (2011). Estrus synchronization in bovine: present status and future prospect. In: *National Symposium on Reproductive Biotechnology for Augmenting Fertility and Conservation of Animal Species with special reference to North Eastern hilly region*, Selesih, Aizwal, Mizoram.
- Perera BMAO (2011). Reproductive cycles of buffalo. *Anim. Reprod. Sci.* 124: 194 – 199.
- Peter AT (2004). An update on cystic ovarian degeneration in cattle. *Reprod. Domest. Anim.* 39: 1 – 7.
- Peter AT, Bosu WTK (1988). Relationship of uterine infections and folliculogenesis in dairy cows during early puerperium. *Theriogenol.* 30: 1045 – 1051.
- Peter AT, Bosu WTK, Gilbert RO (1990). Absorption of *Escherichia coli* endotoxin (lipopolysaccharide) from the uteri of postpartum dairy cows. *Theriogenol.* 33: 189 – 194.
- Peter AT, Vos PLAM, Ambrose DJ (2009). Postpartum anestrus in dairy cattle. *Theriogenol.* 71: 1333 – 1342.
- Peters AR (1984). Effect of exogenous oestradiol-178 on gonadotrophin secretion in post-partum beef cows. *J. Reprod. Fert.* 72: 473 – 478.
- Peters AR, Pimental MG, Lamming GE (1985). Hormone responses to exogenous GnRH pulses in postpartum dairy cows. *J. Reprod. Fert.* 75: 557 – 565.
- Pettersson KJ, Gustafsson H, Strandberg E, Berglund B (2006). Atypical progesterone profiles and fertility in Swedish dairy cows. *J. Dairy Sci.* 89: 2529 – 2538.
- Plasse D, Warnick AC, Koger M (1970). Reproductive behaviour of *Bos indicus* females in a sub-tropical environment. IV. Length of oestrus cycle, duration of oestrus, time of ovulation, fertilization and embryo survival in grade Brahman heifers. *J. Anim. Sci.* 30: 63 – 72.
- Porwal ML, Saxena NK, Shrivastava AM, Karandikar GW (1976). Efficacy of different medicaments on anoestrus buffaloes. *Indian Vet. J.* 53: 453 – 437.
- Prahalad P, Rao KS, Raju KGS (2010). Effect of GnRH, PMSG and Placentex on reproductive performance of postpartum true anoestrus Murrah buffaloes. *Indian J. Anim. Res.* 44(3): 201 – 204.
- Prajapati SB, Ghodasara DJ, Joshi BP, Prajapati KS, Jani VR (2005). Etiopathological study of endometritis in repeat breeder buffaloes. *Buffalo J.* 2: 145 – 165.
- Pryce JE, Coffey MP, Brotherstone S (2000). The genetic relationship between calving interval, body condition score and linear type and management traits in registered Holsteins. *J. Dairy Sci.* 83: 2664.
- Purbey LN, Sane CR (1978). Studies on oestrous cycles in Dangi breed of cows. *Indian Vet. J.* 55: 532 – 535.
- Pursley JR, Mee MO, Wiltbank MC (1995). Synchronization of ovulation in dairy cows using PGF_{2α} and GnRH. *Theriogenol.* 44: 915 – 923.
- Quintana G, Vázquez AI, Weigel KA (2009). Effect of suckling restriction with nose plates and premature weaning on postpartum anoestrus interval in primiparous cows under range conditions. *Anim. Reprod. Sci.* 116: 10 – 18.
- Rahawy MA (2009). Treatment of anoestrus in buffalo cows by massaging the uterus and ovaries rectally. *Iraqi J. Vet. Sci.* 23(1): 23 – 25.
- Rahman MS, Shohag AS, Kamal MM, Parveen N, Shamsuddin M (2012). Application of Ultrasonography to Investigate Postpartum Anestrus in Water Buffaloes. *Reprod Dev Biol* 36(2): 103 – 108.
- Rajkumar R, Srivastava SK, Varshney VP, Mahmood S (2008). Effect of medicinal plants *Saraca asoca* and *Trigonella foenum-graecum* in anoestrus cows. *Indian Vet. J.* 85(12): 1281 – 1283.
- Rao RA (1981). Synchronization of estrus in buffaloes with a progesterone releasing intra-vaginal device. *Vet. Rec.* 159: 123.
- Ramoun AA, Serur BH, El-SM Fattouh, Darweish SA, Abou El-Ghait HA (2012). Enhancing follicular growth as a prerequisite for GnRH treatment of true anoestrus in buffalo. *Anim. Reprod. Sci.* 132: 29 – 35.

- Randel RD (2005). Reproduction of *Bos indicus* breeds and crosses. Proceedings, Applied Reproductive Strategies in Beef Cattle. Texas A&M University, College Station.
- Rao AV, Sreemannarayana O (1982). Clinical analysis of reproductive failure among female buffaloes (*Bubalus bubalis*) under village management in Andhra Pradesh. *Theriogenol.* 18:403 – 411.
- Rao AVN, Reddy AC, Murti YN (1993). Incidence of reproductive disorders in relation to exotic inheritance in cross-bred cows. *Indian Vet. J.* 70: 722 – 725.
- Reddy KRC, Rao AS, Reddy VSC, Yadagiri B, Sharma GP, Reddy MR, Reddy CE (1994). Efficacy of certain hormonal and non-hormonal drugs on estrous induction in postpartum anestrus buffaloes. *Indian J. Anim. Reprod.* 15(2): 127 – 130.
- Rehman S, Wani GM, Khan ZA (1990). Incidence of reproductive disorders in cattle of Kashmir valley. *IJAR.* 11(2) 159.
- Rhodes FM, McDougall S, Burke CR, Verkerk GA, Macmillan KL (2003). Invited Review: Treatment of Cows with an Extended Postpartum Anestrus Interval. *J. Dairy Sci.* 86: 1876 – 1894.
- Rhodes RC, Randel RD (1978). Reproductive studies of Brahman cattle. I. Behavioral effect of various dose levels of estradiol-17 β upon ovariectomized Brahman, Brahman x Hereford and Hereford cows. *Theriogenol.* 9: 429.
- Ribeiro HFL, Andrade de VJ, Marques Jr AP, Vale WG (1997). Effect of body condition at parturition on the interval to first postpartum estrus in buffaloes. *R. Bras. Vet. Med.* 19: 213 – 218.
- Richardson MW, Wettemann RP, Schoenemann HM (1889). Nutritional anestrus in beef cows: concentrations of glucose and non-esterified fatty acids in plasma and insulin in serum. *J. Anim. Sci.* 67: 2354 – 2362.
- Robinson JJ (1990). Nutrition in the reproduction of farm animals. *Nutr. Res. Rev.* 3: 267 – 268.
- Roche JF (2006). The effect of nutritional management of the dairy cow on reproductive efficiency. *Anim. Reprod. Sci.* 96: 282 – 296.
- Roche JF, Mihn M, Diskin MG, Ireland JJ (1998). A review on regulation of follicle growth in cattle. *J. Anim. Sci.* 76 (Suppl. 3): 16 – 19.
- Romaniuk J (1973). Treatment of ovarian afuction in cows. *Medycyna Vet.* 29: 296 – 298.
- Rottensten K, Touchberry RW (1957). Observations on the degree of expression of estrus in cattle. *J. Dairy Sci.* 40: 1457 – 65.
- Roxstrom A, Strandberg E, Berglund B, Emanuelson U, Philipsson J (2001). Genetic and environmental correlations among female fertility traits, and between the ability to show oestrus and milk production in dairy cattle. *Acta. Agric. Scand.* 51: 192 – 199.
- Rutter LM, Randel RD (1984). Postpartum nutrient intake and body condition: effect on pituitary function and onset of oestrus in beef cattle. *J. Anim. Sci.* 58: 265 – 274.
- Ryot KD, Sharma BK, Panwar CD (1990). Effect of iodine therapy in anoestrus bovines. *Indian J. Anim. Reprod.* 11(2): 144 – 145.
- Sah SK, Nakao T (2010). A clinical study of anestrus buffaloes in Southern Nepal. *J. Reprod. Dev.* 56(2): 208 – 211.
- Saiduddin S, Quevedo MM, Foote WD (1968). Response of beef cows to exogenous progesterone and estradiol at various stages postpartum. *J. Anim. Sci.* 27: 1015 – 1020.
- Samad HA, Nasseri AA (1979). A quantitative study of primordial follicles in buffalo heifer ovaries. In: 13th FAO/RSIDA International Course on Animal Reproduction, Uppsala, Sweden.
- Sanchez JM (1995). Iodine in bovine nutrition. *Nutr. Anim. Tropic.* 2: 95 – 120.
- Sane CR, Deshpande BR, Kaikini AS (1967). Problems of infertility in cattle and buffaloes. *Proc. of ICAR Conf. Poona.*
- Sangsrivavong S, Combs DK, Sartori R, Wiltbank MC (2002). High feed intake increases blood flow and metabolism of progesterone and estradiol-17 β in dairy cattle. *J. Dairy Sci.* 85: 2831 – 2842.
- Sarkar AK (2005). Treatment of anestrus cow with diluted logol's iodine and massage on reproductive organs-uncontrolled case study. *J. Anim. Vet. Adv.* 4(8): 734 – 736.
- Selvaraju M, Veerapandian C, Kathiresan D, Chandrasenan C (2005). Incidence of bovine reproductive disorders. *Indian Vet. J.* 82: 556.
- Selvaraju S, Reddy IJ, Gowda NKS, Prasad CS, Ananthram K, Sampath KT (2009). Effect of supplementation of area specific mineral mixture in improving reproductive efficiency in crossbred dairy cattle—a field study. *Indian J. Anim. Sci.* 79 (6): 599 – 601.
- Settergren I (1964). The number of primordial follicles in clinically normal and hypoplastic heifer ovaries, 5th international congress on animal reproduction, Trento, Section IV, 188.
- Settergren I (1997). Ovarian hypoplasia in heifers due to germ cell weakness. *Theriogenol.* 47:31 – 47:539.
- Shah NH, Willemsse AH, Van deWeil DFM, Engel B (1989). Influence of season and parity on several reproductive parameters in Nili-Ravi buffaloes in Pakistan. *Anim. Reprod. Sci.* 21: 177 – 190
- Shah SNH (1990). Prolonged calving intervals in Nilli-Ravi buffalo. PhD dissertation, Rijksuniversiteit Utrecht, pp 162.
- Shams ZH, Kharche KG, Thakur MS (1991). The efficacy of gonadotrophin releasing hormone (Receptal) alone and in combination for oestrus induction in anoestrus crossbred cows. *Indian J. Anim. Reprod.* 12: 175.
- Sharma U, Bhadwal MS, Kumar S (2004). Freemartinism in a crossbred heifer. *Intas Polivet* 5: 164 – 165.
- Sheldon IM, Dobson H (2004). Postpartum uterine health in cattle. *Anim. Reprod. Sci.* 82–83: 295 – 306.
- Short RE, Bellows RA, Staigmiller RB, Berardinelli JG, Custer EE (1990). Physiological mechanisms controlling anestrus and infertility in postpartum beef cattle. *J. Anim. Sci.* 68: 799 – 816.
- Shukla SN, Agarwal SK, Shanker U, Varshney VP, Majumdar AC (2005a). Ovarian function and restoration of fertility using insulin in acyclic dairy cattle. *Indian J. Anim. Sci.* 75: 1135 – 1139.
- Shukla SN, Agarwal SK, Shanker U, Varshney VP, Majumdar AC (2005b). Modulation of ovarian response in anoestrus cattle treated with insulin alone and in combination with GnRH. *Indian J. Anim. Reprod.* 26(2): 159 – 164.
- Silva JRV, Figueiredo JR, Van den Hurk R (2009). Involvement of growth hormone (GH) and insulin-like growth factor (IGF) system in ovarian folliculogenesis. *Theriogenol.* 71: 1193 – 1208.
- Singh CSP, Singh SK, Singh B (1981). Studies on the incidence of infertility in cows. *Indian Vet. J.* 58: 909 – 912.
- Singh G, Dhaliwal GS, Sharma RD, Biswas RK (1988). Treatment of summer anoestrus buffalo (*bubalis bubalis*) with progesterone releasing device plus pregnant mare serum gonadotrophins. *Theriogenol.* 29(5): 1201 – 1206.
- Singh G, Sidhu SS, Verma HK (2003). Incidence of reproductive disorders of buffaloes in different zones of Punjab State. *J. Res.* 40: 79 – 81.
- Singh J, Nanda AS, Adams GP (2000). The reproductive pattern and efficiency of female buffaloes. *Anim. Reprod. Sci.* 60: 593 – 604.
- Singh J, Verma HK, Singh KB, Singh N (2006). Incidence of reproductive disorders in dairy animals in different agroclimatic regions in Punjab. *J. Res.* 43 (3): 224 – 227.
- Singh M, Chaudhari BK, Singh JK, Singh AK, Maurya PK (2013). Effects of thermal load on buffalo reproductive performance during summer season. *J. Biol. Sci.* 1(1): 1 – 8.
- Singh M, Sood P, Vasistha NK, Singh C (2001). Study on the use of prostaglandin F2 α in treatment of suboestrus cows. *Indian Vet. J.* 78: 815 – 816.
- Singh SL, Thakur MS (1999). Estrus synchronization and fertility with the uterine infusion of prostaglandin F2 alpha and lugol's iodine in crossbred cows. *Indian J. Anim. Reprod.* 20(1): 12 – 14.
- Singh V, Malik RK, Singh P, Tuly RK, Verma AK, Chandola RK (2010). Induction of cyclicity in murrh buffaloes—Heifers during summer using different hormonal protocols. *Indian J. Anim. Reprod.* 31(2): 11 – 14.
- Singh VK, Verma HK (1994). Analysis of reproductive disorders of buffaloes. *Liv. Adv.* 19(1): 20 – 21.
- Singhal SP, Dhanda OP, Razdan MN (1984). Some managerial and therapeutic approaches in the treatment of physiological infertility of Water Buffaloes (*Bubalus bubalis*). In: Proceedings of 10th International Congress Animal Reproduction Artificial Insemination. 3: 471.
- Sinha BP, Sinha SN, Singh B (1987). Incidence of anestrus in crossbred cattle in field and farm conditions. *Livestock Advisor.* 12(32): 43 – 48.
- Sirmour S, Nema SP, Singh BK, Shukla SP (2006). Induction of estrus in delayed pubertal crossbred heifer. *Indian J. Anim. Reprod.* 27(1): 55 – 58.
- Sonwane SD, Pargaonkar DR, Bakshi SA, Navtare RM (1995). Studied on efficacy of Receptal to induce estrus in anestrus cow. *Indian J. Anim. Reprod.* 16(2): 132.
- Sonwane SD, Pargaonkar DR, Bakshi SA, Navtake RM, Thakre NV (1994). Studies on serum cholesterol levels on anoestrus cows treated with GnRH analogue (Receptal). *Indian J. Anim. Reprod.* 15(2): 122 – 124.
- Spicer LJ, Convey EM, Tucker HA, Echterkamp SE (1986). Effects of intermittent injections of LHRH on secretory patterns of LH and FSH and ovarian follicular growth during postpartum anovulation in suckled beef cows. *J. Anim. Sci.* 62: 1317 – 1323.
- Stagg K, Spicer LJ, Sreenan JM, Roche JF, Diskin MG (1998). Effect of calf isolation on follicular wave dynamics, gonadotropin and metabolic hormone changes, and interval to first ovulation in beef cows fed either of two energy levels postpartum. *Biol. Reprod.* 59: 777 – 783.
- Reddy SCV, Sharma GP, Raju S, Reddy EC (1990). Effect of clofret-vet treatment in postpartum anoestrus crossbred cows and Murrh Buffaloes. *Indian J. Anim. Reprod.* 11(1): 75 – 76.
- Tapas SP, Shrivastava OP, Pandit RK, Agarwal RG (2000). Estrus response and fertility in sub estrus buffaloes treated with intra uterine medication during low breeding season. *Indian J. Anim. Reprod.* 21(1): 13 – 15.

- Tauk SA, Olsen JR, Wilkinson JRC, Berardinelli JG (2010). Duration of daily bull exposure on resumption of ovulatory activity in postpartum, primiparous, suckled, beef cows. *Anim. Reprod. Sci.* 118: 13 – 18.
- Thakor D, Patel D (2013). Incidence of Infertility Problems in Cattle and Buffaloes. *Dairy Cattle*. <http://en.engormix.com>.
- Thatcher WW, Macmillan KL, Hansen PJ, Drost M (1989). Concepts for regulation of corpus luteum function by the conceptus and ovarian follicles to improve fertility. *Theriogenol.* 31: 149 – 164.
- Tomar DS (2004). Studies on etiology and treatment of subestrus in Murrah buffaloes. M.V.Sc and A.H., Thesis, JNKVV, Jabalpur.
- Ulberg LC, Lindley CE (1960). Use of progesterone and estrogen in the control of reproductive activities in beef cattle. *J. Anim. Sci.* 19: 1132 – 1142.
- Vanholder T, Leroy JLMR, Van Soom A, Opsomer G, Maes D, Kruif (2005). Effect of non-esterified fatty acids on bovine granulosa cell steroidogenesis and proliferation in vitro. *Anim. Reprod. Sci.* 87: 33 – 44.
- Verma HK, Sidhu SS, Panqwanker GR, Dhabilia DC (1992) Treatment of summer anoestrus in buffaloes with Bromocriptine. *Indian J. Anim. Reprod.* 13: 190 – 192.
- Walker SL, Smith RF, Routly JE, Jones DN, Morris MJ, Dobson H (2008). Lameness, activity time–budgets and estrus expression in dairy cattle. *J. Dairy Sci.* 91: 4552 – 4559.
- Walsh RB, Kelton DF, Duffield TF, Leslie KE, Walton JS, LeBlanc SJ (2007). Prevalence and risk factors for postpartum anovulatory condition in dairy cows. *J. Dairy Sci.* 90: 315 – 324.
- Warnick AC, Casida LE, Grummer RH (1950). The occurrence of estrus and ovulation in postpartum sows. *J. Anim. Sci.* 9: 66.
- Webb R, Campbell BK, Garverick HA, Gong JG, Gutierrez CG, Armstrong DG (1999). Molecular mechanisms regulating follicular recruitment and selection. *J. Reprod. Fert. Suppl.* 54: 33 – 48.
- Wettemann RP, Beck TW, Turman EJ, Hintz RL (1982). Endocrine response of postpartum anestrous beef cows to GnRH or PMSG. *Theriogenol.* 18: 599 – 613.
- Whitman RW (1975). Weight change, body condition and beef–cow reproduction. Ph.D. Dissertation. Colorado State Univ., Fort Collins.
- Willet EL (1950). The fertility of heifers following administration of progesterone to alter the estrual cycle. *J. Dairy Sci.* 33: 381
- Williams EJ, Fischer DP, Noakes DE, England GC, Rycroft A, Dobson H (2007). The relationship between uterine pathogen growth density and ovarian function in the postpartum dairy cow. *Theriogenol.* 68: 549 – 559.
- Williams GL (1990). Suckling as a regulator of post–partum rebreeding in cattle: a review. *J. Anim. Sci.* 68: 831 – 852.
- Williams GL, Griffith MK (1995). Sensory and behavioral control of gonadotrophin secretion during suckling–mediated anovulation in cows. *J. Reprod. Fert. Suppl.* 49: 463 – 475.
- Wiltbank JN (1966). Modification of ovarian activity in the bovines following injection of oestrogen and gonadotrophin. *J. Reprod. Fert. Suppl.* 1: 1.
- Wiltbank JN, Cook AC (1958). The comparative reproductive performance of nursed and milked cows. *J. Anim. Sci.* 17: 640 – 648.
- Wiltbank JN, Rowden WW, Ingalls JE, Gregory KE, Koch RM (1962). Effect of energy level on reproductive phenomena of mature Hereford cows. *J. Anim. Sci.* 21: 219.
- Wiltbank JN, Shumway RP, Parker WR, Zimmerman DR (1967). Duration of estrus, time of ovulation and fertilization rate in beef heifer synchronized with DHAP. *J. Anim. Sci.* 26(4): 764.
- Wiltbank M, Gumen A, Sartori R (2002). Physiological classification of anovulatory conditions in cattle. *Theriogenol.* 57: 21 – 52.
- Wordinger RJ, Dickey JF, Hill JR (1976). Influence of a progestogen on carbohydrate histochemical and histologic features of the ampulla of the bovine uterine tube. *Am. J. Vet. Res.* 37: 901.
- Wulji T, Litherland A, Goetsch AL, Sahlou T, Ruchala R, Dawson LJ, Gipson T (2003). Evaluation of melatonin and bromocriptine administration in Spanish goats: effects on the out of season breeding performance in spring, kidding rate and fleece weight of does. *Small Ruminant Res.* 49: 31 – 40.
- Yadav YP, Singh AP, Kunj V, Akhtar MH, Roy GP, Singh C (2004). Study on incidence of anestrous and blood biochemical constituents in non–cyclic and cyclic crossbred cows. *Indian J. Anim. Reprod.* 25: 116 – 119.
- Zakari AY, Molokwu ECI, Osori DIK (1981). Effect of season on the estrus cycle of cows (*Bos indicus* indigenous for northern Nigeria). *Vet. Rec.* 109: 213 – 215.
- Zduńczyk S, Żebracki A, Glazer T, Janowski T, Raś A (1992). Badania nad występowaniem i leczeniem nieczynności jajników u krów w warunkach chowu wielkostatdnego. *Acta Acad Agricult Technol Veterinaria* 20: 87 – 94.
- Zemjanis R, Fahning ML, Schultz RH (1969). Anestrus. *The Practitioner's Dilemma, Vet. Scope (Upjohn)*, 14(1): 14.