

Short Communication

Isolation of *Salmonella* from Liquid Whole Eggs Sold in Retail Outlets in Egypt, Bangladesh and not in Japan

Amira Ahmed^{1,2*}, Hajime Fukunaga¹, Takuya Mizuno¹, Takayuki Ezaki¹

¹Department of Microbiology, Gifu University Graduate School of Medicine, 1-1 Yanagido, Gifu 501-1194, Japan; ²Department of Poultry & Rabbit Medicine, Faculty of Veterinary Medicine, Suez Canal University, Egypt
*Corresponding author: A.A.H.Abdelaziz@gmail.com

ARTICLE HISTORY

Received: 2014-07-27
Revised: 2014-08-07
Accepted: 2014-08-08

ABSTRACT

The main objective of this survey was to establish the current prevalence of *Salmonella* in raw egg samples from the following countries: Japan, Egypt, and Bangladesh. A total of 141 sample units of eggs were tested over a five-month period from December 2013 to April 2014. Following conventional bacteriological procedures the samples subjected for molecular identification by PCR using genus specific primers for *Salmonella* spp. All samples were randomly purchased and analyzed within their stated shelf life. *Salmonella* SPP. contamination was recorded in 17.6% (9/51) of eggs' contents of Egypt samples and recorded in 38% (8/21) of eggs' contents of Bangladesh samples while none of the samples from Japan (0/69) contaminated with *Salmonella* SPP in the eggs' contents. Detection of *Salmonella* in chicken eggs' contents needs a greater concern for effective control of Salmonellosis in poultry.

Key Words: *Salmonella*, Contamination, Poultry, Eggs contents

All copyrights reserved to Nexus® academic publishers

ARTICLE CITATION: Ahmed A; Fukunaga H; Mizuno T; Ezaki T (2014). Isolation of *salmonella* from liquid whole eggs sold in retail outlets in Egypt, Bangladesh and not in Japan. Adv. Anim. Vet. Sci. 2 (7): 390 – 392.

The consumption of raw or lightly cooked eggs, egg containing foods in particular mayonnaise, desserts and sauces has been associated with high percentage of many human Salmonellosis outbreaks.

The intestinal tract is the primary reservoir of *Salmonella* in poultry birds leading to contamination of chicken eggs in cloacal region through horizontal route. Transovarian contamination is another important route of *Salmonella* egg contamination. Several studies have isolated *Salmonella* spp. from ovaries and oviducts of naturally and experimentally infected hens. The presence of *Salmonella* in the reproductive tract was consistent with the production of *Salmonella* contaminated eggs in the albumen, the yolk or the both (Keller et al., 1995; Gast and Beard, 1990; Gast, 1993; Timoney et al., 1989). Food borne *Salmonella* outbreaks still represents a great hazard in Egypt (Ibrahim et al., 2013), Bangladesh (Barua et al., 2013) and in a lower extent in Japan (Suzuki H, and Yamamoto S, 2009).

The goal of this study was to determine the incidence of *Salmonella* in raw egg samples sold in the following countries: Japan, Egypt, and Bangladesh, using both conventional method, and a specific PCR detection technique.

Eggs were purchased from a retail outlets in a residential area of (Gifu Prefecture, Japan; Kafr Elsheik, Dkhalia, and Ismailia cities, Egypt; Dhaka, Bangladesh) over a period of 5-months (December 2013–April 2014). A total of 141 eggs' content were analyzed for the presence of *Salmonella*. In order to collect the eggs' contents, eggs' surfaces were sterilized by swabbing with 70% alcohol, drying with the air and then cracked with a sterile knife. Yolks and whites

were mixed thoroughly, so that their contents pooled; each pool consisted between 6 and 10 eggs.

According to The World Organization for Animal Health 2008, the conventional culture methods for detection of *Salmonella* in food include non-selective pre-enrichment followed by selective enrichment, plating on selective and differential agars, and biochemical tests.

A 25 gram portion of the pooled sample was added to 225 ml BPW and mixed well before incubation at 35° C for 24 hours.

1ml samples were taken and mixed with 10 ml of Rappaport Vassialidis broth (RV) and incubated at 35° C for 24 hours.

A loopful of each enriched sample was streaked on selective agar medium deoxycholate hydrogen sulphide lactose agar (DHL) plate (EIKEN, Tokyo, Japan), using such medium enables testing of the bacterium for lactose fermentation and H₂S production. Suspect colonies are confirmed by molecular identification.

The DNA was released by heating at 100° C for 5 min. The cellular debris was pelleted by centrifugation at 8000 rpm for 5 min, and the clear supernatant fluid containing nucleic acids was conserved until it was used in subsequent PCR assays.

The DNA samples (5 µL) were amplified in an optimized 20 µL reaction mixture consisting of 2.5 µL of each primer 0.4 µM /reaction mixture, (10 µL) 2× SYBER Premix Extaq

buffer (TAKARA, Japan). The reaction mixture was incubated in a programmable DNA thermal cycler (Thermogen, model QB-0224B). *Salmonella* Genus specific primers 139 bp (AMR No DCR227, Japan), which were based on the *invA* gene of *Salmonella*, were used in the PCR assay.

A reagent blank containing all the components of the reaction mixture with the exception of template DNA (which was replaced by sterile distilled water) was included with every PCR assay. Furthermore, positive DNA control was included, which was prepared from *Salmonella* sp., respectively. The cycling parameters used were initially denaturated at 95° C for 1 min followed by 40 cycles of amplification of 10 s at 95° C, and 10 s at 60° C. Then, the PCR tubes were held at 4° C.

The PCR products were analyzed by gel electrophoresis. Five microliters of each sample was loaded onto 2.0% of agarose gel in 1×TAE buffer at 100 V/cm for

30 min. The gel was stained with ethidium bromide and electrophoresed products were visualized with a UV transilluminator (FAS-III, Toyobo, Osaka, Japan). A 100–bp ladder (Toyobo, Japan) was used as a molecular weight marker.




Figure 1: Shows the growth of *Salmonella* on DHL plates with its whitish (non-lactose fermenter) colonies, and blackish colouration because of H₂S production

Growth of *Salmonella* was detected on DHL agar plates as shown in Figure 1 whitish colonies (non-lactose fermenter) with H₂S production indicated by a blackening of the colonies due to a formation of iron sulfide.

Molecular characterization using PCR produced positive amplification of 139 bp fragments of *invA* genes (100%) specific for all members of *Salmonella* SPP.

From the previous results, it could be concluded that *Salmonella*-specific PCR test (Figure 2) in conjunction with traditional isolation methods could be effective in providing a more accurate profile of the prevalence of *Salmonella* in poultry products.

The results of *Salmonella* incidence in eggs from pooled samples are shown in Table 1. *Salmonella* contamination was recorded 17.6% (9/51) of eggs contents of Egypt samples and recorded in 38% (8/21) of eggs contents of Bangladesh

samples while none of the samples from Japan (0/69) contains *Salmonella* contamination in the egg's contents.




Figure 2: A. PCR products of 139 bp specific to *invA* gene of *Salmonella* in ethidium bromide stained 2% agarose TAE gel electrophoresis; M: 100bp; L1: control negative; L2: control positive; L3: PCR product of the sample from Bangladesh pooled eggs; B. PCR products of 139 bp specific to *invA* gene of *Salmonella* in ethidium bromide stained 2% agarose TAE gel electrophoresis. M: 100bp; L1: control negative; L2: control positive; L3: PCR product of the sample from Egypt pooled eggs

Table 1: The prevalence of *Salmonella* isolated from eggs' content

Country of origin	No. of eggs samples	No. of positive samples	of positive samples %
Japan	69	0	0
Egypt	51	9	17.6
Bangladesh	21	8	38

Since *Salmonella* is an important zoonotic bacterium with poultry as largest single reservoir of *Salmonella* (Gupta et al., 1999), the assessment of the contamination level and site of contamination are the utmost importance in deciding the control strategies against *Salmonella* in chickens.

Moreover, the use of a three step protocol (pre-enrichment, selective enrichment and selective plating) as specified in the FDA's Bacteriological Analytical Manual (US Food Drug Administration, 2002) was found satisfactory for the recovery of *Salmonella* spp.

The present study demonstrated widespread contamination by *Salmonella* in eggs retailed in Egypt, similar average of contamination of poultry and or poultry products showed previously by (Ibrahim et al., 2013, Ibrahim et al., 2014)

Our results reinforce previous investigations, suggesting that *Salmonella* as one of the predominant pathogens circulating in poultry in Bangladesh (Barua et al., 2012,2013) and plays a role as a source of human infections (Barua et al., 2014), and consistent with the previous studies indicating the very low internal eggs' contamination in Japan (Suzuki H, and Yamamoto S, 2009).

This study emphasized the need to implement proactive measures of hygienic practices, surveillance programs for laying hen flocks should be optimized, and the application of Hazard Analysis and Critical Control Point (HACCP) in the preparation and processing of foods in

order to reduce *Salmonella* contamination of eggs in retail and to reduce the risk of human infection.

REFERENCES

- Barua H, Biswas PK, Olsen KEP, Christensen JP (2012). Prevalence and characterization of motile *Salmonella* in commercial layer poultry farms in Bangladesh. PLoS ONE 7, e35914, <http://dx.doi.org/10.1371/journal.pone.0035914>.
- Barua H, Biswas PK, Olsen KEP, Shil SK, Christensen JP (2013). Molecular characterization of motile serovars of *Salmonella enterica* from breeder and commercial broiler poultry farms in Bangladesh. PLoS ONE 8, e57811, <http://dx.doi.org/10.1371/journal.pone.0057811>.
- Barua H, Biswas PK, Talukder KA, Olsen KE, Christensen JP (2014). Poultry as a possible source of non-typhoidal *Salmonella enterica* serovars in humans in Bangladesh. Vet. Microbiol. 168: 372 – 380.
- Gast RK, Beard CW (1990). Production of *Salmonella* Enteritidis-contaminated eggs by experimentally infected hens. Avian Dis. 34:438-46.
- Gast RK (1993). Detection of *Salmonella* Enteritidis in experimentally infected laying hens by culturing pools of egg contents. Poul. Sci. 72: 267 – 74.
- Gupta V, Ray P, Sharma M (1999). Antimicrobial resistance pattern of *Shigella* and non-typhi *Salmonella* isolated from patients with diarrhoea. Indian J. Med. Res. 109: 43-45.
- Ibrahim MA, Emeash HH, Ghoneim NH, Abdel-Halim MA (2013). Seroepidemiological studies on poultry *Salmonellosis* and its public health importance. J. World's Poult. Res. 3(1): 18 – 23.
- Ibrahim WA, Abd El-Ghany WA, Nasef SA, Hatem ME (2014). A comparative study on the use of real time polymerase chain reaction (RT-PCR) and standard isolation techniques for the detection of *Salmonellae* in broiler chicks. Int. J. of Vet. Sci. Med. 2 (1): 67 – 71.
- Keller LH, Benson CE, Krotec K, Eckroade RJ (1995). *Salmonella* enteritidis colonization of the reproductive tract and forming and freshly laid eggs of chickens. Infect. Immun. 63: 2443 – 9.
- Suzuki Hodaka, Yamamoto Shigeki (2009). Literature survey of *Salmonella* contamination in eggs and egg products in the world Bulletin of National Institute of Health Sciences. 127: 74 – 83.
- Timoney JF, Shivaprasad HL, Baker RC, Rowe B (1989). Egg transmission after infection of hens with *Salmonella* enteritidis phage type 4. Vet. Rec. 125: 600 – 1.
- US Food Drug Administration (2002). Bacteriological analytical manual. Chap. 5. *Salmonella* AOAC International, Gaithersburg Md Available at <http://www.cfsan.felagovebam/bam-5.html>.