



Supplementary Material

Inoculum Doses and Exposure Periods Affect Recovery of *Steinernema feltiae* and *Heterorhabditis bacteriophora* from *Tenebrio molitor*

Ali Murad Rahoo^{1,*}, Tariq Mukhtar², Ali Murad Jakhar³, Rehana Kanwal Rahoo³

¹Wheat Research Institute, Sakrand, Sindh, Pakistan

²Department of Plant Pathology, Pir Mehr Ali Shah Arid Agriculture University, Rawalpindi, Pakistan

³Institute of Plant Sciences, University of Sindh, Jamshoro, Pakistan

* Corresponding author: alirahoo@gmail.com

0030-9923/2018/0003-0983 \$ 9.00/0

Copyright 2018 Zoological Society of Pakistan

Supplementary Table I.- Analysis of variance.

Source of variation	Degree of freedom	Sum of squares	Mean squares	v. r	F. Pr
Replication	2	1.53	0.76	0.59	
Dose	2	300.96	150.48	116.09	<.001**
Days	5	24.741	4.948	3.82	0.004*
Species	1	27.227	27.227	21.00	<.001**
Dose × Days	10	20.700	2.070	1.60	0.126*
Dose × Species	2	15.520	7.760	5.99	0.004*
Days × Species	5	4.356	0.871	0.67	0.646 ^{NS}
Dose × Days × Species	10	41.233	4.123	3.18	0.002*
Residual	70	90.739	1.296		
Total	107	527.000			

** highly significant; * significant; NS, non-significant.