



Supplementary Material


Short Communication: Prevalence of Foodborne Pathogens in Food Items in Quetta, Pakistan

Abdul Samad¹, Ferhat Abbas¹, Zafar Ahmad¹, Olena Pokryrshko² and Tauseef M Asmat^{1,*}

¹Center for Advanced Studies in Vaccinology and Biotechnology, University of Balochistan, Quetta, Pakistan

²Department of Microbiology, Virology and Immunology, Ternopil State Medical University, Ukraine

* Corresponding author: tauseefcasvab@gmail.com
0030-9923/2018/0004-1597 \$ 9.00/0
Copyright 2018 Zoological Society of Pakistan



Supplementary Fig. S1. Agarose gel (2%) electrophoresis showing amplification fragments of genes of interest. **A**, *invA* genes (284 bp) of *Salmonella* spp. Lanes 1, 2, 6, 8, 10 and 11 show the positive amplification; **B**, *mapA* genes (589 bp) of *Campylobacter jejuni*. Lanes 1, 2, 3, 5, 6, 7, 8 and 9 highlight the positive amplification; **C**, *fliC_{H7}* (625 bp) genes of *E. coli* O157:H7 by Multiplex PCR. Lanes 2, 5, 10 and 12 show the positive amplification; **D**, *hly* genes (730 bp) of *Listeria monocytogenes*. Lanes 1, 2, 4, 7 and 8 denote the positive amplification. PC: Positive Control, NC: Negative Control, M: Molecular weight markers.