



Supplementary Material

Rodents (Muridae) Abundance and Habitat Shift between Agricultural, Non-Agricultural Land and Human Dwellings: A Proposed Strategy to Decrease Crop Damage

Raja Imran Hussain* and Iqra Yousaf

Department of Biology, PMAS-Arid Agriculture University, Rawalpindi

* Corresponding author: rajaimran2003@gmail.com

0030-9923/2019/0006-2051 \$ 9.00/0

Copyright 2019 Zoological Society of Pakistan

Supplementary Table I.- Description of climate characteristics along with altitudinal gradient of study area (Islamabad, Pakistan).

GPS coordinate	Altitude (ASL)	Season	Mean temp. (°C)	Mean precipitation (mm)	Mean monthly sunshine (h)
33.6844° N, 73.0479° E	507~514 m	Autumn	17.25	63.75	268.85
		Spring	12.45	75.80	227.35
		Summer	22.80	169.57	281.77
		Winter	4.65	46.175	206.57

ASL, above sea level.

Supplementary Table II.- Percentage damage caused by rodents to wheat crop during two consecutive harvesting periods (2014-2015).

Crop	Year	Mean number of tillers		Total	Damage (%)
		Whole	Infested		
Wheat	2014	92	34	126	26.98
	2015	66	19	85	22.35