



Research Article

Influence of Biochar based Organic Fertilizers on Growth and Concentration of Heavy Metals in Tomato and Lettuce in Chromite Mine Tailings Contaminated Soil

Faiza Altaf¹, Shamim Gul^{1,2*}, Tasawar Ali Chandio³, Gul Bano Rehman¹, Attiq-ur-Rehman Kakar⁴, Sami Ullah⁴, Naqeebullah Khan⁴, Umbreen Shaheen⁵, Muhammad Naeem Shahwani⁶, Muhammad Ajmal⁷ and Misbah Manzoor⁸

¹Department of Botany, University of Balochistan, Quetta, Pakistan; ²Department of Natural Resource Sciences, McGill University, Quebec, Canada; ³Geological Survey of Pakistan, Saryab road, Quetta, Pakistan; ⁴Department of Chemistry, University of Balochistan, Quetta, Pakistan; ⁵Department of Zoology, University of Balochistan, Quetta, Pakistan; ⁶Faculty of Life Sciences, Balochistan University of Information Technology, Engineering and Management Sciences, Airport Road, 87300, Quetta, Pakistan; ⁷Pakistan Council of Science and Industrial Research, Mastung Road, 87300, Quetta, Balochistan, Pakistan; ⁸Department of Plant Sciences, Sardar Bahadur Khan Women's University, Quetta, Pakistan.


Received | February 29, 2020; Accepted | January 20, 2021; Published | March 08, 2021

*Correspondence | Shamim Gul, Department of Botany, University of Balochistan, Quetta, Pakistan; Email: shamim.gul@mail.mcgill.ca

Citation | Altaf, F., S. Gul, T.A. Chandio, G.B. Rehman, A.u.R. Kakar, S. Ullah, N. Khan, U. Shaheen, M.N. Shahwani, M. Ajmal and M. Manzoor. 2021. Influence of biochar based organic fertilizers on growth and concentration of heavy metals in tomato and lettuce in chromite mine tailings contaminated soil. *Sarhad Journal of Agriculture*, 37(1): 315-324.

DOI | <http://dx.doi.org/10.17582/journal.sja/2021/37.1.315.324>

Keywords | Chromite mine-tailing debris-contaminated soil, Heavy metals, Farmyard manure, Lettuce, Poultry manure, Tomato, Wood-derived biochar



Supplementary Figure S1. Pictures showing debris collected from chromite mines for experiment and plants showing only one fruit per plant or no evident fruit on plants (as only three pots produced only one fruit per plant).



Supplementary Figure S2: Seeds of Roma tomato used in this study.