



Supplementary Materials

Production of Single Cell Microbial Protein and its Use as Protein Source in Broiler Ration

Faheem Ahmed Khan^{1*}, Sarzamin Khan², Qurat-ul-Ain³, Saqib Ishaq¹,
Muhammad Salman¹, Abdul Rehman¹, Ikram Ullah⁴, Kalsoom⁵ and Johar Jamil⁵

¹Department of Microbiology and Biotechnology, Abasyn University, Peshawar, Khyber Pakhtunkhwa, Pakistan

²Department of Poultry Sciences, Faculty of Animal Husbandry, Agricultural University, Peshawar, Khyber Pakhtunkhwa, Pakistan

³Department of Pharmacy, Abasyn University Peshawar, Khyber Pakhtunkhwa, Pakistan

⁴Department of Medical Laboratory Technology, University of Haripur, Khyber Pakhtunkhwa, Pakistan

⁵Department of Microbiology, University of Swabi, Khyber Pakhtunkhwa, Pakistan

Supplementary Table I. Feed formulation (g/kg) of experimental broiler finisher rations with different levels of single cell microbial protein.

S. No.	Ingredients (g/kg)	Group A	Group B	Group C	Group D
1	Soybean meal	68	66	64	70
2	SCMP	2	4	6	0
3	Wheat	205	205	205	205
4	Maize	370	370	370	370
5	Corn gluten meal	48.2	48.2	48.2	48.2
6	Broken rice	50	50	50	50
7	Canola meal	53	53	53	53
8	Guar meal	20	20	20	20
9	Fish meal	70	70	70	70
10	Maize gluten feed	23	23	23	23
11	Molasses	28	28	28	28
12	Rice polishing	43	43	43	43
13	L-Lysine	0.8	0.8	0.8	0.8
14	DL-Methionine	1	1	1	1
15	DCP (di-calcium phosphate)	10	10	10	10
16	Vit. minerals premix	8	8	8	8
	Total	1000	1000	1000	1000

Feed formulation was isonitrogenic and according to NRC 1994.

* Corresponding author: faheemdurani@gmail.com
0030-9923/2021/0003-0875 \$ 9.00/0
Copyright 2021 Zoological Society of Pakistan