



## Supplementary Material


# Short Communication: Epidemiology of Gastrointestinal Parasitism in Blue Wildebeest (*Connochaetes taurinus*), Alpacas (*Vicugna pacos*), and Goats (*Capra aegagrus hircus*) with same Husbandry and Fence Site in Harbin Zoo, China

Yanqiang Zhou<sup>1</sup>, Lixin Wang<sup>1</sup>, Chunxiao Hao<sup>1</sup>, Xiuyun Li<sup>2</sup>, Shakeel Hussain<sup>1</sup>, Dongdong Shen<sup>1</sup>, Zhiwei Peng<sup>1</sup>, Qi'an Zhai<sup>1</sup> and Zhijun Hou<sup>1,\*</sup>

<sup>1</sup>College of Wildlife and Protected Area, Northeast Forestry University, Hexing Road No. 26, Harbin 150040, P.R. China

<sup>2</sup>Harbin Zoo, Harbin, Heilongjiang, P.R. China

Yanqiang Zhou and Lixin Wang equally contributed to this work.



Supplementary Fig. 1. The morphology of oocysts from the three ruminant hosts. A, *Capillaria* sp. in blue wildebeest; B, *Trichuris* sp. in alpacas; C, Strongy-type species in alpacas; D, *Nematodirus* sp. in alpacas; E, *Eimeria macusaniensis* in alpacas; F, *Eimeria* sp I in alpacas; G, *E. aljevi* in goats; H, Strongy-type species in goats; I, *Trichuris* sp. in goats; J, *E. christensenii* in goats.

\* Corresponding author: [houzhihundz@163.com](mailto:houzhihundz@163.com)  
0030-9923/2021/0006-2511 \$ 9.00/0  
Copyright 2021 Zoological Society of Pakistan