




## Supplementary Material

# Diagnosis of Canine Multicentric Lymphoma in Dog


Yuan Zhang<sup>1</sup>, Qingyue Han<sup>1</sup>, Peiquan Du<sup>1</sup>, Yi Lu<sup>1</sup>, Lianmei Hu<sup>1</sup>, Sadaqat Ali<sup>2</sup>, Khalid Mehmood<sup>2</sup>, Sajid Hameed<sup>2</sup>, Zhaoxin Tang<sup>1</sup>, Hui Zhang<sup>1\*</sup> and Ying Li<sup>1\*</sup>

<sup>1</sup>College of Veterinary Medicine, South China Agricultural University, Guangzhou 510642, China.

<sup>2</sup>Faculty of Veterinary and Animal Sciences, Islamia University of Bahawalpur, Bahawalpur, Pakistan.



Supplementary Fig. S1. Thoracic and abdominal X-ray and ultrasonographic images showing pleural effusion (A) and typical nodular lesions in the spleen (B and C).



Supplementary Fig. S2. The histopathological examination of lymphoma in dogs (A), H and E staining for lymphoma cell; the cytological appearance of an immunocytochemical staining for CD3 (B) and CD79a (C) on a lymph node aspirate demonstrating CD79a-reactivity for the neoplastic B-cells and CD3-reactivity for the reactive T-cells consistent with a high-grade B-cell lymphoma). The tumor cells were large and showed a pleomorphic appearance, and the boundaries between the cells were obvious and showed a patchy pattern.

\* Corresponding author: [lying@scau.edu.cn](mailto:lying@scau.edu.cn); [hz236@scau.edu.cn](mailto:hz236@scau.edu.cn)

0030-9923/2022/0002-0957 \$ 9.00/0

Copyright 2022 Zoological Society of Pakistan