



## Supplementary Material

# Solid Waste Management: Impact of Organic Waste on Growth and Reproduction of Earthworm *Eisenia fetida* via Vermicomposting

Irsa Shafique<sup>1</sup>, Saiqa Andleeb<sup>1\*</sup>, Farrukh Naeem<sup>2</sup>, Shaukat Ali<sup>3</sup>,  
Tauseef Tabassum<sup>4</sup> and Tariq Sultan<sup>4</sup>

<sup>1</sup>Microbial Biotechnology and Vermi-technology Laboratory, Department of Zoology, University of Azad Jammu and Kashmir, Muzaffarabad, 13100, Pakistan

<sup>2</sup>First Biotech LLC, Lahore

<sup>3</sup>Department of Zoology, Government College University, Lahore, Pakistan

<sup>4</sup>Land Resources Research Institute, Pakistan Agricultural Research Council, Islamabad, Pakistan




Fig. 1. Vermi-bed preparation and vermi-composting process. (A) Pre-digested fruit and vegetable waste, (B-C) Bedding preparation, (D) Introduction of worms, (E) Introduction of fruit and vegetable waste, (F) Mature clitellate worm, (G) Mixing of material, (H) Introduction of spinach as green material, (I-J) Processing of vermi-compost, (K) Worm movements, (L) Watering, (M-N) Sieving of vermicompost.

\* Corresponding author: [drsaiqa@gmail.com](mailto:drsaiqa@gmail.com), [drsaiqa@ajku.edu.pk](mailto:drsaiqa@ajku.edu.pk)  
0030-9923/2025/0001-0307 \$ 9.00/00



Copyright 2025 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access  article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).