



## Supplementary Material

# Drivers of Electric Pylon Used as Nesting Sites by Birds in Baicheng City, China


Zheng Han<sup>1</sup>, Junbo Liu<sup>2</sup>, Jingyao Luan<sup>2</sup>, Changlong Gao<sup>2</sup>, Yufeng Tai<sup>2</sup>, He Liu<sup>2</sup>, Saipeng Zhang<sup>2</sup>, Guanqiang Zhai<sup>2</sup>, Xi Yang<sup>1,3</sup>, Haitao Wang<sup>1,4\*</sup>

<sup>1</sup>School of Life Sciences, Northeast Normal University, Changchun 130024, China.

<sup>2</sup>Jilin Electric Power Research Institute Co., LTD, Changchun 130021, China.

<sup>3</sup>College of Agricultural, Hulunbuir University, Hulunbuir 021000, China.

<sup>4</sup>Jilin Engineering Laboratory for Avian Ecology and Conservation Genetics, Northeast Normal University, Changchun 130024, China.



Supplementary Fig. S1. The results of spatial autocorrelation in the residuals of the random forest model. The y-axis represents Moran's I estimate, and the x-axis contains the values of the distance thresholds. The dot sizes represent the p-values of the Moran's I estimate, and the black dashed line represents the theoretical null value of the Moran's I estimate. The spatial autocorrelation is weak and non-significant for neighborhood distances larger than 300m.

\* Corresponding author: [wanght402@nenu.edu.cn](mailto:wanght402@nenu.edu.cn)  
0030-9923/2025/0002-0631 \$ 9.00/00



Copyright 2025 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).